

Abundances of mixotrophic, phototrophic, and heterotrophic nanoflagellates, and associated environmental data, synthesized from the scientific literature.

Website: <https://www.bco-dmo.org/dataset/807195>

Data Type: Other Field Results

Version: 1

Version Date: 2020-03-30

Project

» [Eating themselves sick? Ecological interactions among a mixotrophic flagellate, its prokaryotic prey, and an ingestible giant virus.](#) (Giant virus ecology)

Contributors	Affiliation	Role
Edwards, Kyle F.	University of Hawaii at Manoa (SOEST)	Principal Investigator, Contact
Soenen, Karen	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Abundances of mixotrophic, phototrophic, and heterotrophic nanoflagellates, and associated environmental data, synthesized from the scientific literature.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:81 E:25 S:-36.4 W:-125

Temporal Extent: 1990-10-04 - 2018-02-01

Methods & Sampling

Data were compiled from 130 experiments in 11 publications that use fluorescently labeled bacteria or beads to estimate the abundance of bacterivorous mixotrophic nanoflagellates (MNF), defined as nanoflagellates with both ingested labeled prey and chlorophyll autofluorescence. These experiments simultaneously counted heterotrophic nanoflagellates (nanoflagellates with no chlorophyll autofluorescence, HNF) and total phototrophic nanoflagellates (all nanoflagellates with chlorophyll autofluorescence, PNF). In some cases researchers also estimated bulk ingestion rate by MNF and HNF, and these data were also extracted.

To ask whether MNF, HNF, and PNF shift in relative abundance across environmental gradients, data on environmental conditions were extracted from the same studies: temperature, chlorophyll-a concentration, nitrate concentration, and bacterial abundance. Although some samples were taken from below the mixed layer, mixed layer Chl-a and nitrate were used as predictors for all analyses, intended as proxies of total productivity or nutrient supply at that location. For two studies Chl-a was not reported and climatological values were used. To capture the light environment, climatological photosynthetically active radiation data (PAR) were used, because *in situ* PAR data were reported in few studies. Monthly climatological mean PAR from the SeaWiFS mission for each location was extracted from the NASA Giovanni portal (<https://giovanni.gsfc.nasa.gov/giovanni/>).

For samples taken within the mixed layer the light environment was characterized as the median mixed layer PAR, defined as $PAR_{in} * e^{-(0.121 * Chl_{mix}^{0.428} * MLD/2)}$, where PAR_{in} is incident PAR (mole photons per square meter per day), Chl_{mix} is mixed layer Chl-a (micrograms per liter), and MLD is mixed layer depth (meters). For samples taken below the mixed layer the light environment was characterized as PAR at the sample depth, defined as $PAR_{in} * e^{-(0.121 * Chl_{mix}^{0.428} * MLD)}$

For four studies mixed layer depth could not be estimated from data in the study and was taken from a climatology (http://www.ifremer.fr/cerweb/deboyer/mlD/Surface_Mixed_Layer_Depth.php).

Data Processing Description

These data were published in: Edwards, K.F., 2019. Mixotrophy in nanoflagellates across environmental gradients in the ocean. *Proceedings of the National Academy of Sciences*, 116(13), pp.6211-6220. Dataset S1.

BCO-DMO processing notes:

- Changed column headers to comply with database requirements
- Removed reference number variable
- Reformatted the data column to iso format: yyyy-mm-dd
- Rounded the columns Nitrate, Chl_a, HNF, MNF, PNF, Bacteria, HNF_ingest, MNF_infest, MLD, PAR are rounded to 0 decimals
- Rounded the columns HNF, MNF, PNF, Bacteria, HNF_ingest, MNF_infest, MLD are rounded to 0 decimals
- The chl_a, longitude and latitude columns are rounded to 2 decimals
- Rounded PAR to 1 decimal

[[table of contents](#) | [back to top](#)]

Data Files

File
mixotrophic_nanoflagellate.csv (Comma Separated Values (.csv), 12.81 KB) MD5:9bd95ce716a974bb4476e66733fefcf8
Primary data file for dataset ID 807195

[[table of contents](#) | [back to top](#)]

Related Publications

Anderson, R., Jürgens, K., & Hansen, P. J. (2017). Mixotrophic Phytoflagellate Bacterivory Field Measurements Strongly Biased by Standard Approaches: A Case Study. *Frontiers in Microbiology*, 8.

doi:[10.3389/fmicb.2017.01398](https://doi.org/10.3389/fmicb.2017.01398)

Related Research

Arenovski, A. L., Lim, E. L., & Caron, D. A. (1995). Mixotrophic nanoplankton in oligotrophic surface waters of the Sargasso Sea may employ phagotrophy to obtain major nutrients. *Journal of Plankton Research*, 17(4), 801-820. doi:[10.1093/plankt/17.4.801](https://doi.org/10.1093/plankt/17.4.801)

doi:[10.1093/plankt/17.4.801](https://doi.org/10.1093/plankt/17.4.801)

Related Research

Christaki, U., Van Wambeke, F., & Dolan, J. (1999). Nanoflagellates (mixotrophs, heterotrophs and autotrophs) in the oligotrophic eastern Mediterranean: standing stocks, bacterivory and relationships with bacterial production. *Marine Ecology Progress Series*, 181, 297-307. doi:[10.3354/meps181297](https://doi.org/10.3354/meps181297)

Related Research

Czypionka, T., Vargas, C. A., Silva, N., Daneri, G., González, H. E., & Iriarte, J. L. (2011). Importance of mixotrophic nanoplankton in Aysén Fjord (Southern Chile) during austral winter. *Continental Shelf Research*,

31(3-4), 216–224. doi:[10.1016/j.csr.2010.06.014](https://doi.org/10.1016/j.csr.2010.06.014)

Related Research

Edwards, K. F. (2019). Mixotrophy in nanoflagellates across environmental gradients in the ocean. *Proceedings of the National Academy of Sciences*, 116(13), 6211–6220. doi:[10.1073/pnas.1814860116](https://doi.org/10.1073/pnas.1814860116)

Results

Gast, R. J., Fay, S. A., & Sanders, R. W. (2018). Mixotrophic Activity and Diversity of Antarctic Marine Protists in Austral Summer. *Frontiers in Marine Science*, 5. doi:[10.3389/fmars.2018.00013](https://doi.org/10.3389/fmars.2018.00013)

Related Research

Moorthi, S., Caron, D., Gast, R., & Sanders, R. (2009). Mixotrophy: a widespread and important ecological strategy for planktonic and sea-ice nanoflagellates in the Ross Sea, Antarctica. *Aquatic Microbial Ecology*, 54, 269–277. doi:[10.3354/ame01276](https://doi.org/10.3354/ame01276)

Related Research

Safi, K., & Hall, J. (1999). Mixotrophic and heterotrophic nanoflagellate grazing in the convergence zone east of New Zealand. *Aquatic Microbial Ecology*, 20, 83–93. doi:[10.3354/ame020083](https://doi.org/10.3354/ame020083)

Related Research

Sanders, R. W., & Gast, R. J. (2011). Bacterivory by phototrophic picoplankton and nanoplankton in Arctic waters. *FEMS Microbiology Ecology*, 82(2), 242–253. doi:[10.1111/j.1574-6941.2011.01253.x](https://doi.org/10.1111/j.1574-6941.2011.01253.x)

Related Research

Sanders, R., Berninger, U., Lim, E., Kemp, P., & Caron, D. (2000). Heterotrophic and mixotrophic nanoplankton predation on picoplankton in the Sargasso Sea and on Georges Bank. *Marine Ecology Progress Series*, 192, 103–118. doi:[10.3354/meps192103](https://doi.org/10.3354/meps192103)

Related Research

Sato, M., Shiozaki, T., & Hashihama, F. (2016). Distribution of mixotrophic nanoflagellates along the latitudinal transect of the central North Pacific. *Journal of Oceanography*, 73(2), 159–168. doi:[10.1007/s10872-016-0393-x](https://doi.org/10.1007/s10872-016-0393-x)

Related Research

Vargas, C., Contreras, P., & Iriarte, J. (2012). Relative importance of phototrophic, heterotrophic, and mixotrophic nanoflagellates in the microbial food web of a river-influenced coastal upwelling area. *Aquatic Microbial Ecology*, 65(3), 233–248. doi:[10.3354/ame01551](https://doi.org/10.3354/ame01551)

Related Research

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Method	fluorescently labeled bacteria (FLB) or fluorescently labeled beads (beads)	unitless
Group	grouping variable for within-study spatial/temporal structure	unitless
Environment	coastal or open ocean (defined by eye)	unitless
Latitude	latitude, negative is south	decimal degrees
Longitude	longitude, negative is west	decimal degrees
Position	mixed layer (ML) or below the mixed layer	unitless
Nitrate	nitrate concentration	micromoles per liter (umol/L)
Chl_a	mixed layer Chlorophyl a	micrograms per liter (ug/L)
HNF	heterotrophic nanoflagellates	cells per milliliter (cells/mL)
MNF	mixotrophic nanoflagellates	cells per milliliter (cells/mL)
PNF	phototrophic nanoflagellates	cells per milliliter (cells/mL)
Bacteria	bacteria	cells per milliliter (cells/mL)
HNF_ingest	total bacteria ingested by heterotrophic nanoflagellates	cells ingested per hour
MNF_ingest	total bacteria ingested by mixotrophic nanoflagellates	cells ingested per hour
MLD	mixed layer depth	meter (m)
PAR	climatological photosynthetically active radiation data (PAR) - either median mixed layer PAR or PAR at sample depth, as defined in Methods	mol photons per m ² per day
Study	publication data are compiled from - full reference can be found in "publications" section of the BCO-DMO dataset landing page	unitless
Date	Date in format YYYY-MM-DD	unitless
Depth	Depth from the surface.	meters (m)
Temperature	Water temperature at sample depth	degrees Celsius (°C)

[[table of contents](#) | [back to top](#)]

Project Information

Eating themselves sick? Ecological interactions among a mixotrophic flagellate, its prokaryotic prey, and an ingestible giant virus. (Giant virus ecology)

Coverage: North Pacific Subtropical Gyre - Station ALOHA; and North Pacific tropical embayment, Oahu, HI - Kaneohe Bay

NSF Award Abstract:

Phytoplankton support the biological bounty of our seas, so understanding what controls their growth and death is one of the central issues in oceanography. In much of the nutrient-depleted surface waters of the open ocean, the most successful phytoplankton are tiny photosynthetic bacteria known as Prochlorococcus. These bacteria are highly successful competitors for the ocean's limited nutrients and commonly outcompete larger phytoplankton. Yet, many larger types of phytoplankton persist in the ocean. One reason why this

coexistence may occur is that some of the weaker competitors called mixotrophs have evolved a clever alternative strategy best summed up as "If you can't beat them, eat them". In addition to directly competing for nutrients dissolved in the water, these larger phytoplankton can acquire nutrients by consuming and digesting their smaller rivals. The dual ability to photosynthesize and eat competitors has clear advantages, but there can be hidden costs of this intraguild predation strategy. While feeding on *Prochlorococcus*, mixotrophs may also inadvertently ingest giant viruses that are so large they are mistaken for food. Infection is often fatal. Mixotrophy and viral infection are ubiquitous in the ocean; however these processes are often understudied and missing from traditional models of marine food webs that generally consider photosynthesis and predation independently. In this project, the interactions among a common mixotroph (*Florenciella*), its prey (*Prochlorococcus*), and a virus that infects the mixotroph (FloV1) will be studied in the lab and field. This research will also help guide the development of a cohesive mixotroph-virus-prey trophic model. Improving these trophic models to account for more complex processes could fundamentally change our understanding of marine trophic dynamics. The project will directly support the training of a post-doc, graduate and undergraduate student in inter-disciplinary science that includes field, lab, and modeling activities. The project will support a major component of the graduate student's dissertation and the progressive training of an undergraduate student, culminating in an independent project. The concepts of mixotrophy and viral ecology investigated here will be translated into a public display seen by hundreds of children and members of the public. The PIs will engage a K-12 teacher in the fieldwork at sea through a "Science Teachers Aboard Research Ships (STARS)" program and will recruit an undergraduate researcher through the CMORE Scholars program at the University of Hawaii.

The advantages and drawbacks of a mixotrophic strategy will depend on the availability of resources and competitors and the likelihood of viral infection. The timing of grazing will be tested to determine whether *Florenciella* grazes continuously or separates its grazing and photosynthetic activities by only feeding at night. Prey preferences of *Florenciella* will be tested in competitive grazing experiments offering *Prochlorococcus* as prey in the presence of varying amounts of other bacteria and cyanobacteria. Electron microscopy will be used to determine whether prey and virus enter *Florenciella* by the same pathway and whether the presence of prey competitively interferes with viral infection. The kinetics of grazing by *Florenciella* and infection of *Florenciella* by FloV1 will be quantified. The results from these lab experiments will be used to parameterize a numerical model. The model will be used to answer questions and make predictions about the dynamics of the mixotroph-virus-prey system and those predictions will be compared to field data. Collectively, these observational, experimental and quantitative analyses will provide a detailed exploration of the ecological complexity hidden at the base of the marine food web.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1559356

[[table of contents](#) | [back to top](#)]