

Minnow trap data obtained in the Corpus Christi Bay and Mission-Aransas Bays, Texas, USA between July and August, 2018.

Website: <https://www.bco-dmo.org/dataset/813563>

Data Type: Other Field Results

Version: 1

Version Date: 2020-05-30

Project

» [RAPID: Degradation and Resilience of Seagrass Ecosystem Structure and Function following a Direct Impact by Hurricane Harvey](#) (Harvey Seagrass)

Contributors	Affiliation	Role
Yeager, Lauren	University of Texas - Marine Science Institute (UTMSI)	Principal Investigator, Contact
Dunton, Kenneth	University of Texas - Marine Science Institute (UTMSI)	Co-Principal Investigator
Soenen, Karen	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Minnow trap data obtained in the Corpus Christi Bay and Mission-Aransas Bays, Texas, USA between July and August, 2018.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:27.9774 E:-97.09258 S:27.188541 W:-97.1544

Temporal Extent: 2018-07-26 - 2018-08-09

Dataset Description

Minnow trap data obtained in the Corpus Christi Bay and Mission-Aransas Bays, Texas, USA between July and August, 2018.

Methods & Sampling

Targeted sampling of seagrass nekton was conducted at 8 sites along seagrass edges and interiors to determine whether seagrass edges may function differently than seagrass interiors as habitat. At each sites, we set 2 pairs of minnow traps. For each pair, one trap was set directly along the seagrass edge and the second was set within seagrass habitat at least 2 m from the edge. Each trap was baited with a small handful of dry dogfood. The traps were left to soak for ~24 hours. Upon collection, fishes and invertebrates were placed into a bucket of seawater, identified and measured to the nearest mm. Each site was sampled once in July and once in August 2018.

Data Processing Description

BCO-DMO processing notes:

- Adjusted column headers to comply with database requirements
- Added ISO datetime (Set + Retrieved datetimes) fields
- Changed , to ; in field column to comply with database requirements

[[table of contents](#) | [back to top](#)]

Data Files

File
nektion.csv (Comma Separated Values (.csv), 91.67 KB) MD5:7f2fb7a2c4b4720e70621df386030e8c Primary data file for dataset ID 813563

[[table of contents](#) | [back to top](#)]

Related Publications

Congdon, V. M., Bonsell, C., Cuddy, M. R., & Dunton, K. H. (2019). In the wake of a major hurricane: Differential effects on early vs. late successional seagrass species. *Limnology and Oceanography Letters*, 4(5), 155–163. doi:[10.1002/lo2.10112](https://doi.org/10.1002/lo2.10112)

Methods

Duffy, J. E., Ziegler, S. L., Campbell, J. E., Bippus, P. M., & Lefcheck, J. S. (2015). Squidpops: A Simple Tool to Crowdsource a Global Map of Marine Predation Intensity. *PLOS ONE*, 10(11), e0142994.

doi:[10.1371/journal.pone.0142994](https://doi.org/10.1371/journal.pone.0142994)

Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Unique_id	Unique trap identifier: Site_replicate_position_month_sampling.gear	unitless
Site	Site name where traps set	unitless
Checked	Y	unitless
Latitude	Latitude, south is negative. Taken at trap with handheld GPS	decimal degrees
Longitude	Longitude, west is negative. Taken at trap with handheld GPS	decimal degrees
Site_rep	Replicate pair at site (1 or 2)	unitless
Trap_label	Unique label on minnow trap, helps with retrieval	unitless
Month	Month full name.	unitless
Date_set	Date (mm-dd-yy) the trap was set in US Central Time	unitless
Date_retrieved	Date (mm-dd-yy) the trap was retrieved in US Central Time	unitless
Time_set	Time (hh:mm) of day the trap was set in US Central Time	unitless

Time_retrieved	Time (hh:mm) of day the trap was retrieved in US Central Time	unitless
Depth_set_cm	Depth where the trap was set, measured with meter stick to closest cm	centimeters (cm)
Position_set	Was the trap set at the edge of interior of a seagrass bed	unitless
Notes	Field notes	unitless
Common_name	Species of fauna in trap	unitless
Scientific_name	Species of fauna in trap	unitless
Total_N	Total abundance for this species	count
L1	Length (SL or CW) of each individual recorded	millimeters (mm)
L2	Length (SL or CW) of each individual recorded	millimeters (mm)
L3	Length (SL or CW) of each individual recorded	millimeters (mm)
L4	Length (SL or CW) of each individual recorded	millimeters (mm)
L5	Length (SL or CW) of each individual recorded	millimeters (mm)
L6	Length (SL or CW) of each individual recorded	millimeters (mm)
L7	Length (SL or CW) of each individual recorded	millimeters (mm)
L8	Length (SL or CW) of each individual recorded	millimeters (mm)
L9	Length (SL or CW) of each individual recorded	millimeters (mm)
L10	Length (SL or CW) of each individual recorded	millimeters (mm)
L11	Length (SL or CW) of each individual recorded	millimeters (mm)
L12	Length (SL or CW) of each individual recorded	millimeters (mm)
L13	Length (SL or CW) of each individual recorded	millimeters (mm)
L14	Length (SL or CW) of each individual recorded	millimeters (mm)
L15	Length (SL or CW) of each individual recorded	millimeters (mm)
L16	Length (SL or CW) of each individual recorded	millimeters (mm)
L17	Length (SL or CW) of each individual recorded	millimeters (mm)
L18	Length (SL or CW) of each individual recorded	millimeters (mm)
L19	Length (SL or CW) of each individual recorded	millimeters (mm)
L20	Length (SL or CW) of each individual recorded	millimeters (mm)

L21	Length (SL or CW) of each individual recorded	millimeters (mm)
L22	Length (SL or CW) of each individual recorded	millimeters (mm)
L23	Length (SL or CW) of each individual recorded	millimeters (mm)
L24	Length (SL or CW) of each individual recorded	millimeters (mm)
L25	Length (SL or CW) of each individual recorded	millimeters (mm)
L26	Length (SL or CW) of each individual recorded	millimeters (mm)
L27	Length (SL or CW) of each individual recorded	millimeters (mm)
L28	Length (SL or CW) of each individual recorded	millimeters (mm)
L29	Length (SL or CW) of each individual recorded	millimeters (mm)
L30	Length (SL or CW) of each individual recorded	millimeters (mm)
L31	Length (SL or CW) of each individual recorded	millimeters (mm)
L32	Length (SL or CW) of each individual recorded	millimeters (mm)
L33	Length (SL or CW) of each individual recorded	millimeters (mm)
L34	Length (SL or CW) of each individual recorded	millimeters (mm)
L35	Length (SL or CW) of each individual recorded	millimeters (mm)
L36	Length (SL or CW) of each individual recorded	millimeters (mm)
L37	Length (SL or CW) of each individual recorded	millimeters (mm)
L38	Length (SL or CW) of each individual recorded	millimeters (mm)
L39	Length (SL or CW) of each individual recorded	millimeters (mm)
L40	Length (SL or CW) of each individual recorded	millimeters (mm)
L41	Length (SL or CW) of each individual recorded	millimeters (mm)
L42	Length (SL or CW) of each individual recorded	millimeters (mm)
L43	Length (SL or CW) of each individual recorded	millimeters (mm)
L44	Length (SL or CW) of each individual recorded	millimeters (mm)
L45	Length (SL or CW) of each individual recorded	millimeters (mm)

L46	Length (SL or CW) of each individual recorded	millimeters (mm)
L47	Length (SL or CW) of each individual recorded	millimeters (mm)
L48	Length (SL or CW) of each individual recorded	millimeters (mm)
L49	Length (SL or CW) of each individual recorded	millimeters (mm)
L50	Length (SL or CW) of each individual recorded	millimeters (mm)
L51	Length (SL or CW) of each individual recorded	millimeters (mm)
L52	Length (SL or CW) of each individual recorded	millimeters (mm)
L53	Length (SL or CW) of each individual recorded	millimeters (mm)
L54	Length (SL or CW) of each individual recorded	millimeters (mm)
L55	Length (SL or CW) of each individual recorded	millimeters (mm)
L56	Length (SL or CW) of each individual recorded	millimeters (mm)
L57	Length (SL or CW) of each individual recorded	millimeters (mm)
L58	Length (SL or CW) of each individual recorded	millimeters (mm)
L59	Length (SL or CW) of each individual recorded	millimeters (mm)
L60	Length (SL or CW) of each individual recorded	millimeters (mm)
L61	Length (SL or CW) of each individual recorded	millimeters (mm)
L62	Length (SL or CW) of each individual recorded	millimeters (mm)
L63	Length (SL or CW) of each individual recorded	millimeters (mm)
L64	Length (SL or CW) of each individual recorded	millimeters (mm)
L65	Length (SL or CW) of each individual recorded	millimeters (mm)
L66	Length (SL or CW) of each individual recorded	millimeters (mm)
L67	Length (SL or CW) of each individual recorded	millimeters (mm)
L68	Length (SL or CW) of each individual recorded	millimeters (mm)
L69	Length (SL or CW) of each individual recorded	millimeters (mm)

L70	Length (SL or CW) of each individual recorded	millimeters (mm)
L71	Length (SL or CW) of each individual recorded	millimeters (mm)
L72	Length (SL or CW) of each individual recorded	millimeters (mm)
L73	Length (SL or CW) of each individual recorded	millimeters (mm)
L74	Length (SL or CW) of each individual recorded	millimeters (mm)
L75	Length (SL or CW) of each individual recorded	millimeters (mm)
L76	Length (SL or CW) of each individual recorded	millimeters (mm)
L77	Length (SL or CW) of each individual recorded	millimeters (mm)
L78	Length (SL or CW) of each individual recorded	millimeters (mm)
L79	Length (SL or CW) of each individual recorded	millimeters (mm)
L80	Length (SL or CW) of each individual recorded	millimeters (mm)
L81	Length (SL or CW) of each individual recorded	millimeters (mm)
L82	Length (SL or CW) of each individual recorded	millimeters (mm)
L83	Length (SL or CW) of each individual recorded	millimeters (mm)
L84	Length (SL or CW) of each individual recorded	millimeters (mm)
L85	Length (SL or CW) of each individual recorded	millimeters (mm)
L86	Length (SL or CW) of each individual recorded	millimeters (mm)
L87	Length (SL or CW) of each individual recorded	millimeters (mm)
L88	Length (SL or CW) of each individual recorded	millimeters (mm)
L89	Length (SL or CW) of each individual recorded	millimeters (mm)
L90	Length (SL or CW) of each individual recorded	millimeters (mm)
L91	Length (SL or CW) of each individual recorded	millimeters (mm)
L92	Length (SL or CW) of each individual recorded	millimeters (mm)
L93	Length (SL or CW) of each individual recorded	millimeters (mm)

L94	Length (SL or CW) of each individual recorded	millimeters (mm)
L95	Length (SL or CW) of each individual recorded	millimeters (mm)
L96	Length (SL or CW) of each individual recorded	millimeters (mm)
L97	Length (SL or CW) of each individual recorded	millimeters (mm)
L98	Length (SL or CW) of each individual recorded	millimeters (mm)
L99	Length (SL or CW) of each individual recorded	millimeters (mm)
L100	Length (SL or CW) of each individual recorded	millimeters (mm)
L101	Length (SL or CW) of each individual recorded	millimeters (mm)
L102	Length (SL or CW) of each individual recorded	millimeters (mm)
L103	Length (SL or CW) of each individual recorded	millimeters (mm)
L104	Length (SL or CW) of each individual recorded	millimeters (mm)
L105	Length (SL or CW) of each individual recorded	millimeters (mm)
L106	Length (SL or CW) of each individual recorded	millimeters (mm)
L107	Length (SL or CW) of each individual recorded	millimeters (mm)
L108	Length (SL or CW) of each individual recorded	millimeters (mm)
L109	Length (SL or CW) of each individual recorded	millimeters (mm)
L110	Length (SL or CW) of each individual recorded	millimeters (mm)
L111	Length (SL or CW) of each individual recorded	millimeters (mm)
L112	Length (SL or CW) of each individual recorded	millimeters (mm)
L113	Length (SL or CW) of each individual recorded	millimeters (mm)
L114	Length (SL or CW) of each individual recorded	millimeters (mm)
L115	Length (SL or CW) of each individual recorded	millimeters (mm)
L116	Length (SL or CW) of each individual recorded	millimeters (mm)
L117	Length (SL or CW) of each individual recorded	millimeters (mm)

L118	Length (SL or CW) of each individual recorded	millimeters (mm)
L119	Length (SL or CW) of each individual recorded	millimeters (mm)
L120	Length (SL or CW) of each individual recorded	millimeters (mm)
L121	Length (SL or CW) of each individual recorded	millimeters (mm)
L122	Length (SL or CW) of each individual recorded	millimeters (mm)
L123	Length (SL or CW) of each individual recorded	millimeters (mm)
L124	Length (SL or CW) of each individual recorded	millimeters (mm)
L125	Length (SL or CW) of each individual recorded	millimeters (mm)
L126	Length (SL or CW) of each individual recorded	millimeters (mm)
L127	Length (SL or CW) of each individual recorded	millimeters (mm)
L128	Length (SL or CW) of each individual recorded	millimeters (mm)
L129	Length (SL or CW) of each individual recorded	millimeters (mm)
L130	Length (SL or CW) of each individual recorded	millimeters (mm)
ISO_DateTime_Set_UTC	Datetime of trap setting in ISO format (YYYY-mm-ddTHH:MMZ) in UTC time zone.	unitless
ISO_DateTime_Retrieved_UTC	Datetime of trap retrieved in ISO format (YYYY-mm-ddTHH:MMZ) in UTC time zone.	unitless

[[table of contents](#) | [back to top](#)]

Project Information

RAPID: Degradation and Resilience of Seagrass Ecosystem Structure and Function following a Direct Impact by Hurricane Harvey (Harvey Seagrass)

Coverage: Corpus Christi Bay and Mission-Aransas Bays, Texas, USA

NSF Award Abstract:

Disturbance has long been recognized as a major organizing force in marine communities with the potential to shape biodiversity. Hurricanes provide a natural experiment to understand how acute physical disturbances (storm surge and wind energy) may interact with longer-term changes in environmental conditions (salinity or turbidity) to alter the structure and function of ecological communities. As models indicate that hurricane intensity and precipitation will increase with a warming climate, understanding the response and recovery of coastal ecosystems is of critical societal importance. Harvey made landfall as a Category Four hurricane on the Texas coast on August 25, 2017, bringing extreme rainfall as the storm stalled over the middle Texas coast. The heavy rainfall and freshwater run-off created a low salinity lens that continues to persist two months later.

Seagrass ecosystems may be particularly vulnerable because they grow on shallow, soft-sediment bottoms (and thus are easily dislodged or buried) and because seagrasses are sensitive to changes in salinity and turbidity. The societal implications of seagrass loss are well recognized: seagrasses provide highly valuable ecosystem services of large economic value for estuarine and nearshore dependent fisheries, serve as nursery habitats, and sequester gigatons of carbon on a global scale. Using measurements of the health and function of the seagrass and of the community for which it is habitat, the PIs are assessing the impact of the hurricane and of the persistent freshwater lens. Context is provided by looking at non-impacted sites and by six prior years of data.

This project addresses the overarching question: How do intense physical disturbances in conjunction with chronic chemophysical perturbations affect loss and recovery of seagrass community structure and function, including local production, trophic linkages, and metazoan community diversity? To understand the impacts of Hurricane Harvey on seagrass ecosystems across the middle Texas coast, the investigators are (1) documenting losses in physical habitat structure, (2) teasing apart independent and interactive effects of multiple stressors associated with storm events on biodiversity and ecosystem function, and (3) identifying factors that promote resilience following disturbance. A state-wide seagrass monitoring program with six years of data from areas within Harvey's path and surrounding seagrass systems will provide invaluable context. The investigators are measuring seagrass structure, employing a Before-After-Control-Impact design at sites that experienced severe physical damage and appropriate reference sites. In situ loggers deployed after the storm track the evolution of the low salinity event together with seagrass physiological stress measurements (e.g. chlorophyll fluorescence, pigment loss, reduced growth). Changes in seagrass habitat function is assessed through measurements of faunal biodiversity within impacted and reference sites sampled via cores, benthic push nets, and seine nets. Tethering assays of seagrass blades and common invertebrate prey enables comparison trophic interactions across sites that vary in disturbance impact. These data are used to create models of ecosystem response to an extreme disturbance event and identify factors that best predict recovery of the physical structure of the habitat and of associated ecosystem functions.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1807143

[[table of contents](#) | [back to top](#)]