

Growth of Antarctic krill (*Euphausia superba*) from a 1-month experiment at ambient conditions, ambient temperature and elevated pCO₂, or elevated temperature and elevated pCO₂, Jan-Feb. 2014

Website: <https://www.bco-dmo.org/dataset/820395>

Data Type: experimental

Version: 1

Version Date: 2020-08-10

Project

» [Collaborative Research: Synergistic effects of Elevated Carbon Dioxide \(CO₂\) and Temperature on the Metabolism, Growth, and Reproduction of Antarctic Krill \(*Euphausia superba*\)](#) (OA Krill)

Contributors	Affiliation	Role
Saba, Grace	Rutgers University	Principal Investigator
Seibel, Brad	University of South Florida (USF)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Antarctic krill (*Euphausia superba*) growth data collected during a 1-month experiment where krill were maintained at either ambient conditions, ambient temperature and elevated pCO₂, or elevated temperature and elevated pCO₂.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [Data Files](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: Lat:-64.7741 Lon:-64.0526

Temporal Extent: 2014-01-18 - 2014-02-26

Dataset Description

Antarctic krill (*Euphausia superba*) growth data collected during a 1-month experiment where krill were maintained at either ambient conditions, ambient temperature and elevated pCO₂, or elevated temperature and elevated pCO₂.

Methods & Sampling

Sampling and analytical procedures:

Capture and husbandry: Antarctic krill (*Euphausia superba*) were captured during the austral summer of 2013/2014. Krill were collected by net tow (2 m diameter, 1000 m mesh, non-filtering cod end) off the R/V Laurence M. Gould near the Western Antarctic Peninsula and transported directly to the Palmer Station

biological laboratory. One to two thousand krill were housed in one 4'w x 3'h circular holding tank and two 5' x 2' x 1' rectangular tanks provided with aeration and flow-through seawater. Water was non-filtered and individuals were able to feed on plankton ad libitum throughout the season.

Experimental treatments: Three experimental treatments were targeted in this study, (1) ambient temperature and ambient pCO₂/pH (400ppm/8.10), (2) ambient temperature and elevated pCO₂ (800ppm)/reduced pH (7.7), and (3) elevated temperature (3 degrees C) and elevated pCO₂ (800ppm)/reduced pH (7.7). Temperature treatments were obtained using two separate recirculating systems. Two 800 L cylindrical polycarbonate carboys were attached to temperature controlled chillers (Delta Star) and inline pumps. The carboys were filled with non-filtered seawater acquired from the Palmer Station intake line, placed in a flow-through water bath, and maintained at 0 degrees C. Another 800 L carboy was set up without a chiller and placed in an environmental chamber set at 3 degrees C. The systems were replaced with new water daily and allowed to acclimate to temperature for a minimum of 24 hours before the start of a trial or water change. High CO₂ conditions were obtained using a peristaltic pump to inject straight CO₂ into the propeller of a pump submerged in seawater. Treated water was then gently siphoned with minimal disturbance into treatment bottles.

Krill of comparable size (23-32 mm) were picked and placed in individual 4 L wide-mouth polycarbonate bottles with airtight lids (n = 30 per experimental treatment). Bottles were filled with one of the three pre-acclimated treatment waters as described above. Bottles were immediately closed and placed in environmental chambers set to either 0 or 3 degrees C. Every 24-48 hours, 80% of the water was siphoned out of each experimental bottle and replaced with pre-acclimated water from the 800 L carboys to minimize excretory and respiratory effect of the animals on treatment conditions. During the 39-day experiment, bottles were checked for molts twice per day. If a molt was observed, that bottle was pulled from the experiment and the molt and live krill processed. The experiment ended after the last krill molted.

Analyses: Once a krill molted, both the krill and molt were gently removed from the bottle. Both the molt and krill were measured for total length (mm) with digital calipers (Fowler), and the krill wet weight (mg) was measured.

Salinity, pH, and total alkalinity were measured for each experimental treatment at the start of the experiment and at each water change (every 24-48 hours) for the duration of the incubation. Salinity was measured with a bench top conductivity meter (YSI 3100) calibrated daily with a conductivity standard (50,000 uS/cm; Ricca Chemical Company). pH was determined spectrophotometrically using the indicator dye thymol blue (Dickson et al. 2007; Zhang and Byrne 1996). Total alkalinity was determined on 100 ml subsamples with an open-cell, potentiometric titration of seawater (Metrohm 888 Titrando) with 0.1 M HCl following the potential of a pH electrode (Dickson et al. 2007). Tiamo software (version 2.3) was used to process the alkalinity data. Measurements of pH and TA were quality controlled using certified reference materials (CRMs) obtained from Andrew Dickson at UCSD Scripps Institute of Oceanography

Data Processing Description

BCO-DMO Processing Notes:

- data submitted in Excel file "2014_AntarcticKrill_GrowthExperiment.xlsx" sheet "Sheet1" extracted to csv
- removed carbonate table to server separately
- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- converted dates to ISO format (yyyy-mm-dd)
- replaced ND with nd for 'no data'

[[table of contents](#) | [back to top](#)]

Data Files

File	
2014_krill_growth.csv	(Comma Separated Values (.csv), 5.94 KB) MD5:8e95229624f7e60c79f6d474854fe186
Primary data file for dataset ID 820395	
Carbonate chemistry data from the 2014 Antarctic krill growth experiments	(Comma Separated Values (.csv), 3.56 KB) MD5:afcce3b829a688384f69e326c13d2b4f
filename: 2014_carbonate.csv	
Temperature, pCO ₂ , pH, total alkalinity in sample bottles for 2014 growth experiments.	
BCO-DMO Processing Notes:	
<ul style="list-style-type: none"> - carbonate data submitted in Excel file "2014_AntarcticKrill_GrowthExperiment.xlsx" sheet "Sheet1" extracted to csv - changed date format to ISO format (yyyy-mm-dd) - modified parameter names to conform with BCO-DMO naming conventions - replaced NA with nd for 'no data' 	

[[table of contents](#) | [back to top](#)]

Related Publications

Dickson, A.G., Sabine, C.L. and Christian, J.R. (Eds.) 2007. Guide to best practices for ocean CO₂ measurements. PICES Special Publication 3, 191 pp. ISBN: 1-897176-07-4. URL: https://www.nodc.noaa.gov/ocads/oceans/Handbook_2007.html <https://hdl.handle.net/11329/249>
Methods

Zhang, H., & Byrne, R. H. (1996). Spectrophotometric pH measurements of surface seawater at in-situ conditions: absorbance and protonation behavior of thymol blue. *Marine Chemistry*, 52(1), 17-25.
doi:[10.1016/0304-4203\(95\)00076-3](https://doi.org/10.1016/0304-4203(95)00076-3)
Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Bottle_Krill_no	bottle number	unitless
Treatment_temperature	Temperature treatment; either Ambient (0 degrees C) or Elevated (3 degrees C)	degrees Celsius
Treatment_pCO ₂	pCO ₂ treatment; either Ambient or Elevated (target = 800 ppm)	unitless
Experiment_Start_Date	Date experiment was begun; formatted as yyyy-mm-dd	unitless
Molt_Date	Date krill molted in bottle; bottle pulled from experiment on this date; formatted as yyyy-mm-dd	unitless
Duration	Length of time between start of experiment and time of molt;	days
Total_krill_length_pre_molt	Krill length determined from the molt	millimeters (mm)
Live_krill_total_length_post_molt	Krill length determined from the live krill after it molted	millimeters (mm)
Krill_wet_weight	Wet weight determined from the live krill after it molted;	mg
Notes	comments on experiment	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Delta Star chiller
Generic Instrument Name	Aquarium chiller
Dataset-specific Description	Used to cool water to ambient temperature.
Generic Instrument Description	Immersible or in-line liquid cooling device, usually with temperature control.

Dataset-specific Instrument Name	Metrohm 888 Titrande
Generic Instrument Name	Automatic titrator
Dataset-specific Description	Instrument used for open-cell titrations to determine total alkalinity in seawater. Used to measure total alkalinity in seawater.
Generic Instrument Description	Instruments that incrementally add quantified aliquots of a reagent to a sample until the end-point of a chemical reaction is reached.

Dataset-specific Instrument Name	Fowler digital calipers
Generic Instrument Name	calipers
Dataset-specific Description	Used to measure krill uropod and total lengths.
Generic Instrument Description	A caliper (or "pair of calipers") is a device used to measure the distance between two opposite sides of an object. Many types of calipers permit reading out a measurement on a ruled scale, a dial, or a digital display.

Dataset-specific Instrument Name	YSI 3100 Conductivity Instrument
Generic Instrument Name	Conductivity Meter
Dataset-specific Description	Used to measure salinity in seawater.
Generic Instrument Description	Conductivity Meter - An electrical conductivity meter (EC meter) measures the electrical conductivity in a solution. Commonly used in hydroponics, aquaculture and freshwater systems to monitor the amount of nutrients, salts or impurities in the water.

Dataset-specific Instrument Name	
Generic Instrument Name	Plankton Net
Dataset-specific Description	Net with 2 m diameter, 1000 m mesh, non-filtering cod end. Used to collected krill for experimental analyses.
Generic Instrument Description	A Plankton Net is a generic term for a sampling net that is used to collect plankton. It is used only when detailed instrument documentation is not available.

Dataset-specific Instrument Name	Shimadzu spectrophotometer
Generic Instrument Name	Spectrophotometer
Dataset-specific Description	Used to measure pH in seawater.
Generic Instrument Description	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Synergistic effects of Elevated Carbon Dioxide (CO₂) and Temperature on the Metabolism, Growth, and Reproduction of Antarctic Krill (*Euphausia superba*) (OA Krill)

Website: <http://coseenow.net/project-parka/>

Coverage: Palmer Station, Antarctica

NSF Award Abstract:

Climate change projections for this century suggest that the Southern Ocean will be the first region to be affected by seawater chemistry changes associated with enhanced carbon dioxide (CO₂). Additionally, regions of the Southern Ocean are warming faster than any other locations on the planet. Ocean acidification and warming may act synergistically to impair the performance of different organisms by simultaneously increasing metabolic needs and reducing oxygen transport. However, no studies have measured krill acid-base regulation, metabolism, growth, or reproduction in the context of ocean acidification or synergistic "greenhouse" conditions of elevated CO₂ and temperature. In the present project, the investigators will conduct both short and prolonged exposure experiments at Palmer Station, Antarctica to determine the responses of *Euphausia superba* to elevated CO₂ and temperature. The investigators will test hypotheses related to acid-base compensation and acclimation of various life stages of krill to elevated CO₂ and temperature. Furthermore, they will determine these impacts on feeding, respiration, metabolism, growth, and reproduction.

The Antarctic krill, *Euphausia superba*, is a key species in Antarctic food webs as they are a primary food source for many of the top predators in the Southern Ocean including baleen whales, seals, penguins, and other sea birds. This project will determine the responses of Antarctic krill exposed to elevated CO₂ and temperature and whether or not krill have the capacity to fully compensate under future ocean conditions. The proposed field effort will be complemented by an extensive broader impact effort focused on bringing marine science to both rural and urban high school students in the Midwest (Kansas). The core educational objectives of this proposal are to 1) instruct students about potential careers in marine science, 2) engage students and promote their interest in the scientific process, critical thinking, and applications of science, mathematics, and technology, and 3) and increase student and teacher awareness and understanding of the oceans and global climate change, with special focus on the Western Antarctic Peninsula region. Finally, this project will engage

undergraduate and graduate students in the production, analysis, presentation and publication of datasets.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Office of Polar Programs (formerly NSF PLR) (NSF OPP)	OPP-1641198
NSF Office of Polar Programs (formerly NSF PLR) (NSF OPP)	OPP-1246293

[[table of contents](#) | [back to top](#)]