

# R code for a model of phytoplankton community structure under multiple frequencies of pulsed nutrient supply.

**Website:** <https://www.bco-dmo.org/dataset/820488>

**Data Type:** model results

**Version:** 1

**Version Date:** 2020-08-11

## Project

» [Eating themselves sick? Ecological interactions among a mixotrophic flagellate, its prokaryotic prey, and an ingestible giant virus.](#) (Giant virus ecology)

Contributors	Affiliation	Role
<a href="#">Edwards, Kyle F.</a>	University of Hawaii at Manoa (SOEST)	Principal Investigator, Contact
<a href="#">Soenen, Karen</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

R code for a model of phytoplankton community structure under multiple frequencies of pulsed nutrient supply.

---

## Table of Contents

- [Dataset Description](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Dataset Description

A trait-based approach is used to model how phytoplankton community structure might vary when nutrients are supplied periodically at multiple timescales.

Trait variation across phytoplankton 'species' is defined by an empirically supported three-way tradeoff between maximum growth rate, specific uptake affinity for phosphate, and internal storage capacity for phosphorus (Edwards et al. 2013).

400 species varying along on this tradeoff plane are initialized in the community, and competition proceeds, with species at very low abundance removed, until the dynamics converge on a periodic attractor.

The frequency and magnitude of nutrient pulses are varied to investigate how community trait structure and diversity respond. Nutrient pulses are represented as mixing events with water below the surface mixed layer, which simultaneously dilute the phytoplankton populations. Variations in parameter values and the resulting changes in community structure are described in Smith and Edwards (2019).

## Data Processing Description

BCO-DMO processing notes:

- Linked to the provided R code

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>Phytoplankton_Community_Structure_Rcode_Smith_Edwards_2019</b> filename: Phytoplankton_Community_Structure_Rcode_Smith_Edwards_2019.R (Octet Stream, 10.34 KB) MD5:08bfe78a634c2d96e6d8d55292db92c2
R code to model changes in phytoplankton community structure and diversity, based on three traits: maximum growth rate, specific uptake affinity for phosphate, and internal storage capacity for phosphorus. These variables are affected by periodically supplying nutrients at multiple timescales.
parameter definitions: umax = maximum growth rate Psaff = scaled uptake affinity for phosphorous vmaxhi = maximum cell-specific phosphate uptake rate when quota is at its minimum vmaxlo = maximum uptake rate when quota is at its maximum K = half-saturation constant for phosphate uptake Q = cellular internal phosphorus concentration Qmin = minimum phosphorus quota $u.infin = umax * Qmax / (Qmax - Qmin)$ = theoretical growth rate at infinite quota m = instantaneous mortality rate a = instantaneous mixing rate d = effective instantaneous mixing/mortality rate for the pulses S = dissolved phosphorus concentration in water below the mixed layer f = proportion of phosphorus in dead cells that is instantly remineralized

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Edwards, K. F., Klausmeier, C. A., & Litchman, E. (2013). A Three-Way Trade-Off Maintains Functional Diversity under Variable Resource Supply. *The American Naturalist*, 182(6), 786–800. doi:[10.1086/673532](https://doi.org/10.1086/673532)  
*Results*

Smith, A. N., & Edwards, K. F. (2019). Effects of multiple timescales of resource supply on the maintenance of species and functional diversity. *Oikos*, 128(8), 1123–1135. doi:[10.1111/oik.04937](https://doi.org/10.1111/oik.04937)  
*Results*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

*Parameters for this dataset have not yet been identified*

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

**Eating themselves sick? Ecological interactions among a mixotrophic flagellate, its prokaryotic prey, and an ingestible giant virus. (Giant virus ecology)**

**Coverage:** North Pacific Subtropical Gyre - Station ALOHA; and North Pacific tropical embayment, Oahu, HI - Kaneohe Bay

### *NSF Award Abstract:*

Phytoplankton support the biological bounty of our seas, so understanding what controls their growth and death is one of the central issues in oceanography. In much of the nutrient-depleted surface waters of the open ocean, the most successful phytoplankton are tiny photosynthetic bacteria known as *Prochlorococcus*. These bacteria are highly successful competitors for the ocean's limited nutrients and commonly outcompete larger phytoplankton. Yet, many larger types of phytoplankton persist in the ocean. One reason why this coexistence may occur is that some of the weaker competitors called mixotrophs have evolved a clever alternative strategy best summed up as "If you can't beat them, eat them". In addition to directly competing for nutrients dissolved in the water, these larger phytoplankton can acquire nutrients by consuming and digesting their smaller rivals. The dual ability to photosynthesize and eat competitors has clear advantages, but there can be hidden costs of this intraguild predation strategy. While feeding on *Prochlorococcus*, mixotrophs may also inadvertently ingest giant viruses that are so large they are mistaken for food. Infection is often fatal. Mixotrophy and viral infection are ubiquitous in the ocean; however these processes are often understudied and missing from traditional models of marine food webs that generally consider photosynthesis and predation independently. In this project, the interactions among a common mixotroph (*Florenciella*), its prey (*Prochlorococcus*), and a virus that infects the mixotroph (FloV1) will be studied in the lab and field. This research will also help guide the development of a cohesive mixotroph-virus-prey trophic model. Improving these trophic models to account for more complex processes could fundamentally change our understanding of marine trophic dynamics. The project will directly support the training of a post-doc, graduate and undergraduate student in inter-disciplinary science that includes field, lab, and modeling activities. The project will support a major component of the graduate student's dissertation and the progressive training of an undergraduate student, culminating in an independent project. The concepts of mixotrophy and viral ecology investigated here will be translated into a public display seen by hundreds of children and members of the public. The PIs will engage a K-12 teacher in the fieldwork at sea through a "Science Teachers Aboard Research Ships (STARS)" program and will recruit an undergraduate researcher through the CMORE Scholars program at the University of Hawaii.

The advantages and drawbacks of a mixotrophic strategy will depend on the availability of resources and competitors and the likelihood of viral infection. The timing of grazing will be tested to determine whether *Florenciella* grazes continuously or separates its grazing and photosynthetic activities by only feeding at night. Prey preferences of *Florenciella* will be tested in competitive grazing experiments offering *Prochlorococcus* as prey in the presence of varying amounts of other bacteria and cyanobacteria. Electron microscopy will be used to determine whether prey and virus enter *Florenciella* by the same pathway and whether the presence of prey competitively interferes with viral infection. The kinetics of grazing by *Florenciella* and infection of *Florenciella* by FloV1 will be quantified. The results from these lab experiments will be used to parameterize a numerical model. The model will be used to answer questions and make predictions about the dynamics of the mixotroph-virus-prey system and those predictions will be compared to field data. Collectively, these observational, experimental and quantitative analyses will provide a detailed exploration of the ecological complexity hidden at the base of the marine food web.

[ [table of contents](#) | [back to top](#) ]

---

## **Funding**

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1559356</a>

[ [table of contents](#) | [back to top](#) ]