

Photosynthetically active radiation (PAR) from Odyssey PAR sensors deployed at three sites in the Florida Keys in June 2018

Website: <https://www.bco-dmo.org/dataset/821036>

Data Type: Other Field Results

Version: 1

Version Date: 2020-08-17

Project

» [Carbon Cycling in Carbonate-Dominated Benthic Ecosystems: Eddy Covariance Hydrogen Ion and Oxygen Fluxes](#) (ECHOES Benthic Ecosystems)

Contributors	Affiliation	Role
Long, Matthew H.	Woods Hole Oceanographic Institution (WHOI)	Principal Investigator
McCorkle, Daniel C.	Woods Hole Oceanographic Institution (WHOI)	Co-Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

An eddy covariance system, known as ECHOES, was deployed at three sites offshore of Key Largo, Florida during June 2018. The ECHOES systems logged the three-dimensional velocity, depth, O₂ optode, pH sensor, and triaxial Inertial Measurement Unit. A separate frame at each site contained a photosynthetically active radiation (PAR) sensor and a Seabird SeapHOx, measuring salinity, temperature, depth, O₂, and pH. This dataset contains the PAR data.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:25.120328 E:-80.300504 S:25.11732 W:-80.303069

Temporal Extent: 2018-06-24 - 2018-06-29

Dataset Description

An eddy covariance system, known as ECHOES, was deployed at three sites offshore of Key Largo, Florida during June 2018. The ECHOES systems logged the three-dimensional velocity, depth, O₂ optode, pH sensor, and triaxial Inertial Measurement Unit. A separate frame at each site contained a photosynthetically active radiation (PAR) sensor and a Seabird SeapHOx, measuring salinity, temperature, depth, O₂, and pH. This dataset contains the PAR data.

Methods & Sampling

Background

The basis for the eddy covariance (EC) technique is that turbulent mixing, caused by the interaction of current velocity with the benthic, atmospheric, sea-ice, or cline interfaces, is the dominant vertical transport process in boundary layers. Therefore, vertical fluxes across the ecosystem interfaces can be derived from high-resolution measurements of the vertical velocity and a solute concentration.

Field Sites

The field sites were located ~7 km offshore of Key Largo, Florida, USA at the southern tip of Florida in the Florida Keys. The sites were located on or adjacent to Little Grecian Rocks Reef with a site on the reef crest (25.119016°N, -80.300504°W) at 2.9 m mean depth, in a seagrass bed located ~225 m to the northwest of the reef site (25.120328°N, -80.302222°W) at 4.8 m mean depth, and in a sandy site located ~300 m to the southwest of the reef site (25.117320°N, -80.303069°W) at 6.3 m mean depth. The reef site is described in substantial detail (3-dimensional and species analyses) in Hopkinson et al. (2020), where the EC instrument can be seen near the center of the image analyses (in Figure 6 of Hopkinson et al. 2020) during its deployment in this study. This reef site is substantially degraded with its benthic surface and primary production dominated by octocorals, algae and rubble (Hopkinson et al. 2020). The seagrass site was dominated by dense *Thalassia testudinum* (turtlegrass) with a canopy height of 0.2 m underlain by carbonate sands. The sandy site was composed of carbonate sands with microalgal mats and migrating bedforms 0.1 m in height. Research was conducted from June 24 to June 29 in 2018 with the seagrass deployment beginning on the 24th and the sand and reef deployment beginning on the 25th of June, 2018.

Instrumentation

The EC systems used here, known as Eddy Covariance Hydrogen Ion and Oxygen Exchange System (ECHOES, Long et al. 2015) consisted of an Acoustic Doppler Velocimeter (ADV, Nortek) that was coupled to a FirestingO₂ Mini fiber-optic O₂ meter with a fast-response (~ 0.3 s) 430 μm diameter optode (Pyroscience) (Long et al. 2015, Long and Nicholson 2018, Long et al. 2019) and a fast-response (~0.6 s) Honeywell Durafet III pH sensor with a preamp Cap Adapter and a custom isolation amplifier (based on Texas Instruments ISO124P). The ECHOES systems logged the three-dimensional velocity, depth, O₂ optode, pH sensor, and triaxial Inertial Measurement Unit (IMU, MicroStrain model 3DM-GX3) at a frequency of 32 Hz continuously.

A separate frame at each site contained an Odyssey (Dataflow Systems, New Zealand) photosynthetically active radiation (PAR) sensor and a Seabird SeapHOx (measuring salinity, temperature, depth, O₂, and pH). The SeapHOx was factory calibrated and the Odyssey PAR sensors were calibrated to a HR-4 spectroradiometer system (HOBI Labs HydroRAD-4) using the methods of Long et al. (2012).

Refer to the Supplemental File "ECHOES_methods_FL2018.pdf" for more details on methodology.

Data Processing Description

BCO-DMO Processing:

- concatenated data from the three different sites into one dataset;
- renamed fields;
- added latitude and longitude columns.

[[table of contents](#) | [back to top](#)]

Data Files

File
Fl_par.csv (Comma Separated Values (.csv), 46.79 KB) MD5:abb8624ba3f689e75dc5bf534986fc20
Primary data file for dataset ID 821036

[[table of contents](#) | [back to top](#)]

Supplemental Files

File

ECHOES FL 2018 Methods

filename: ECHOES_methods_FL2018.pdf

(Portable Document Format (.pdf), 615.40 KB)
MD5:32e714361f0a5aee33baae62748d4333

Study site and methods description for Florida Keys 2018 datasets from the project "Carbon Cycling in Carbonate-Dominated Benthic Ecosystems: Eddy Covariance Hydrogen Ion and Oxygen Fluxes".

[[table of contents](#) | [back to top](#)]

Related Publications

Hopkinson, B. M., King, A. C., Owen, D. P., Johnson-Roberson, M., Long, M. H., & Bhandarkar, S. M. (2020). Automated classification of three-dimensional reconstructions of coral reefs using convolutional neural networks. PLOS ONE, 15(3), e0230671. doi:[10.1371/journal.pone.0230671](https://doi.org/10.1371/journal.pone.0230671)

Methods

Long, M. H., & Nicholson, D. P. (2017). Surface gas exchange determined from an aquatic eddy covariance floating platform. Limnology and Oceanography: Methods, 16(3), 145–159. doi:[10.1002/lom3.10233](https://doi.org/10.1002/lom3.10233)

Methods

Long, M. H., Charette, M. A., Martin, W. R., & McCorkle, D. C. (2015). Oxygen metabolism and pH in coastal ecosystems: Eddy Covariance Hydrogen ion and Oxygen Exchange System (ECHOES). Limnology and Oceanography: Methods, 13(8), 438–450. doi:[10.1002/lom3.10038](https://doi.org/10.1002/lom3.10038)

Methods

Long, M. H., Koopmans, D., Berg, P., Rysgaard, S., Glud, R. N., & Søgaard, D. H. (2012). Oxygen exchange and ice melt measured at the ice-water interface by eddy correlation. Biogeosciences, 9(6), 1957–1967.

doi:[10.5194/bg-9-1957-2012](https://doi.org/10.5194/bg-9-1957-2012)

Methods

Long, M. H., Rheuban, J. E., McCorkle, D. C., Burdige, D. J., & Zimmerman, R. C. (2019). Closing the oxygen mass balance in shallow coastal ecosystems. Limnology and Oceanography, 64(6), 2694–2708.

doi:[10.1002/lno.11248](https://doi.org/10.1002/lno.11248)

Methods

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
site	Site description	unitless
lat	Site latitude	decimal degrees North
lon	Site longitude	decimal degrees East
PAR_time	hour (from 00:00 on 6/24/2018)	hours
Photosynthetically_Active_Radiation	Photosynthetically Active Radiation (PAR)	milli Einsteins per square meter per second (mE m ⁻² s ⁻¹)

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Odyssey (Dataflow Systems, New Zealand) photosynthetically active radiation (PAR) sensor
Generic Instrument Name	Photosynthetically Available Radiation Sensor
Generic Instrument Description	A PAR sensor measures photosynthetically available (or active) radiation. The sensor measures photon flux density (photons per second per square meter) within the visible wavelength range (typically 400 to 700 nanometers). PAR gives an indication of the total energy available to plants for photosynthesis. This instrument name is used when specific type, make and model are not known.

Dataset-specific Instrument Name	HR-4 spectroradiometer system (HOBI Labs HydroRAD-4)
Generic Instrument Name	Radiometer
Generic Instrument Description	Radiometer is a generic term for a range of instruments used to measure electromagnetic radiation (radiance and irradiance) in the atmosphere or the water column. For example, this instrument category includes free-fall spectral radiometer (SPMR/SMSR System, Satlantic, Inc), profiling or deck cosine PAR units (PUV-500 and 510, Biospherical Instruments, Inc). This is a generic term used when specific type, make and model were not specified.

[[table of contents](#) | [back to top](#)]

Project Information

Carbon Cycling in Carbonate-Dominated Benthic Ecosystems: Eddy Covariance Hydrogen Ion and Oxygen Fluxes (ECHOES Benthic Ecosystems)

Website: <https://www2.whoi.edu/staff/mlong/projects/project-4/>

Coverage: Bermuda

NSF Award Abstract:

Chemical and biological processes that occur in and on the seafloor can create chemical exchange of elements with seawater and make significant contributions to carbon and nutrient cycling in shallow coastal systems. However, these processes are exceedingly difficult to measure directly in the ocean, with no satisfactory methods currently available to quantify their full impact. The researchers undertaking this project have developed a unique, field instrument referred to as the Eddy Covariance H⁺ and O₂ Exchange System (ECHOES). These novel measurements of hydrogen ion (H⁺) and oxygen (O₂) exchange between the seafloor and the overlying seawater will allow unique, direct evaluation of the important linked biological and chemical reactions. Data from ECHOES will transform understanding of the potentially critical contribution of seafloor processes to the resilience of coastal ecosystems experiencing rapid changes in seawater chemistry. Results from this project will provide critical data for improved models of the consequences of coastal acidification. Additionally, this project will fund an early career scientist and the mentorship of undergraduate students in ocean science research through the Woods Hole Oceanographic Institute's Summer Student Fellowship Program.

Laboratory experiments have successfully examined the benthic response of individual organisms and chemical reactions to stress related to changing seawater chemistry but the integrated response of intact ecosystems has been very difficult to quantify due to unsatisfactory methods for in situ measurements of the required suite of biogeochemical fluxes. This deployment of ECHOES at a variety of carbonate-dominated

seafloor sites in Bermuda is a pioneering effort to simultaneously measure net community production (NCP) and net community calcification (NCC). The study will focus on traditionally difficult-to-study systems including complex reefs, vertical seagrass canopies, and bare permeable sediments, evaluating diel variability, patchiness, and the impact of upstream fluxes on downstream ecosystems. Important biogeochemical parameters (e.g. pH, CO₂, O₂, alkalinity, etc.) in these productive shallow environments can experience daily fluctuations over a greater dynamic range than 100-year model projections for the open ocean due to increasing atmospheric CO₂. Therefore, the novel field data generated by this research will help define the potentially critical and heretofore ill-defined role for shallow, productive carbonate sediments in predictive models of ecosystem response to ocean acidification.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1657727

[[table of contents](#) | [back to top](#)]