Surface pCO2, bottle-corrected ChI a, and shipboard wind speed from cruises conducted during 2010, 2011, and 2012 off the Northeast coast of Brazil in the Amazon River Plume as part of the ANACONDAS project

Website: https://www.bco-dmo.org/dataset/849870

Data Type: Cruise Results

Version: 1

Version Date: 2021-04-20

Project

» Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses (ANACONDAS)

Programs

- » Marine Microbiology Initiative (MMI)
- » Integrated Marine Biogeochemistry and Ecosystem Research US (IMBER-US)
- » Ocean Carbon and Biogeochemistry (OCB)
- » Emerging Topics in Biogeochemical Cycles (ETBC)

Contributors	Affiliation	Role
Yager, Patricia L.	University of Georgia (UGA)	Principal Investigator
Mu, Linquan	University of Georgia (UGA)	Contact
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Surface pCO2, bottle-corrected ChI a, and shipboard wind speed from three cruises conducted off the Northeast coast of Brazil in the Amazon River Plume as part of the ANACONDAS project. Cruise AN10 took place on R/V Knorr (KN197-08) from May-June 2010; cruise AN11 took place on R/V Melville (MV1110) from September-October 2011; and cruise AN12 took place on R/V Atlantis (AT21-04) during July 2012.

Table of Contents

- Coverage
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- Related Publications
- <u>Parameters</u>
- <u>Instruments</u>
- Deployments
- <u>Project Information</u>
- <u>Program Information</u>
- Funding

Coverage

Spatial Extent: N:13.1 **E**:-44.64 **S**:-0.2166 **W**:-59.63

Temporal Extent: 2010-06-06 - 2012-07-29

Methods & Sampling

Methodology Summary

SST and SSS were continuously monitored using an in-line Seabird thermosalinograph. Chlorophyll a

fluorescence (FLR) was recorded by a fluorometer mounted 5-7 m below sea surface, which was then converted to chlorophyll a concentration (Chla) using a linear regression of FLR against discrete Chla measurements of surface water samples collected with Niskin bottles during each cruise. True wind speed was measured shipboard by meteorological instruments on each vessel. Wind speed at 10 m above sea surface (U10) was corrected from instrument heights based on Thomas et al. (2005). Partial pressure of carbon dioxide at sea surface (pCO2_sw) was determined with a showerhead-type pCO2 analyzer (Bates et al., 1998; Sweeney et al., 2000; Takahashi et al., 1997) and an infrared CO2 detector at in-situ SST (Cooley et al., 2007).

pCO2 sampling procedures

The volumetric concentration of CO_2 (xCO_2 (sw), in ppm) was automatically calibrated at least hourly against six CO_2 - N_2 gas mixtures containing 0–510 ppm CO_2 (Certified by the National Oceanographic and Atmospheric Administration Climate Monitoring and Diagnostics Laboratory, NOAA CMDL, Boulder, Colorado, USA). Sea surface pCO_2 was converted from xCO_2 (sw) and corrected to in-situ SST (see Weiss & Price, 1980, and Takahashi et al., 1993, for details). The gas stream between the equilibrator and the detector was dried, but this filter sometimes failed, leaving gaps in the data along the cruise track where water vapor overwhelmed infrared detection of CO_2 . These system failures were apparent and easily removed manually. Instrument accuracy was $\pm 1~\mu$ atm and the response (e-folding; Cooley et al., 2007) time was 6, 2, 10 mins for cruises during June 2010, July 2012, and September 2011, respectively.

More details on methodology can be found in Mu et al. (2021).

Data Processing Description

BCO-DMO Processing:

- concatenated separate data sheets into one dataset;
- added cruise id field:
- converted "Time" field to ISO DateTime UTC.

[table of contents | back to top]

Data Files

File

AN10-12_pCO2_chla.csv(Comma Separated Values (.csv), 1.01 MB) MD5:8b89525775ad61cebcd6841051724051

Primary data file for dataset ID 849870

[table of contents | back to top]

Related Publications

Bates, N. R., Takahashi, T., Chipman, D. W., & Knap, A. H. (1998). Variability of pCO2 on diel to seasonal timescales in the Sargasso Sea near Bermuda. Journal of Geophysical Research: Oceans, 103(C8), 15567–15585. doi:10.1029/98jc00247 https://doi.org/10.1029/98jC00247 Methods

Cooley, S. R., & Yager, P. L. (2006). Physical and biological contributions to the western tropical North Atlantic Ocean carbon sink formed by the Amazon River plume. Journal of Geophysical Research, 111(C8). doi:10.1029/2005jc002954 https://doi.org/10.1029/2005JC002954 Methods

Mu, L., Gomes, H. do R., Burns, S. M., Goes, J. I., Coles, V. J., Rezende, C. E., Thompson, F. L., Moura, R. L., Page, B., & Yager, P. L. (2021). Temporal Variability of Air-Sea CO2 flux in the Western Tropical North Atlantic Influenced by the Amazon River Plume. Global Biogeochemical Cycles, 35(6). Portico. https://doi.org/10.1029/2020gb006798 https://doi.org/10.1029/2020gb006798 https://doi.org/10.1029/2020GB006798

Sweeney, C., Hansell, D. A., Carlson, C. A., Codispoti, L. ., Gordon, L. I., Marra, J., ... Takahashi, T. (2000).

Biogeochemical regimes, net community production and carbon export in the Ross Sea, Antarctica. Deep Sea Research Part II: Topical Studies in Oceanography, 47(15-16), 3369–3394. doi:10.1016/s0967-0645(00)00072-2 https://doi.org/10.1016/S0967-0645(00)00072-2 Methods

Takahashi, T., Feely, R. A., Weiss, R. F., Wanninkhof, R. H., Chipman, D. W., Sutherland, S. C., & Takahashi, T. T. (1997). Global air-sea flux of CO2: An estimate based on measurements of sea-air pCO2 difference. Proceedings of the National Academy of Sciences, 94(16), 8292–8299. doi:10.1073/pnas.94.16.8292 Methods

Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., & Sutherland, S. C. (1993). Seasonal variation of CO2and nutrients in the high-latitude surface oceans: A comparative study. Global Biogeochemical Cycles, 7(4), 843–878. doi:10.1029/93gb02263 https://doi.org/10.1029/93GB02263 *Methods*

Thomas, B. R., Kent, E. C., & Swail, V. R. (2005). Methods to homogenize wind speeds from ships and buoys. International Journal of Climatology, 25(7), 979–995. doi: 10.1002/joc.1176
Methods

Weiss, R. F., & Price, B. A. (1980). Nitrous oxide solubility in water and seawater. Marine Chemistry, 8(4), 347-359. doi: 10.1016/0304-4203(80)90024-9 General

[table of contents | back to top]

Parameters

Parameter	Description	Units
cruise_name	ANACONDAS project cruise name	unitless
cruise_id	Official cruise identifier	unitless
ISO_DateTime_UTC	Date and time (UTC) formatted to ISO8601 standard: YYYY-MM-DDThh:mm:ssZ	unitless
Time	Ordinal date & time (within a year)	unitless
Latitude	Latitude	decimal degrees North
Longitude	Longitude	decimal degrees East
SST	Sea surface temperature	degrees Celsius
SSS	Sea surface salinity	unitless
Chl_a	Chlorophyll a concentration	milligrams per liter (mg L-1)
U10	Wind speed at 10 m height	meters per second (m s-1)
pCO2sw	Surface water pCO2	microatmospheres (uatm)

[table of contents | back to top]

Instruments

Dataset- specific Instrument Name	
Generic Instrument Name	Fluorometer
Generic Instrument	A fluorometer or fluorimeter is a device used to measure parameters of fluorescence: its intensity and wavelength distribution of emission spectrum after excitation by a certain spectrum of light. The instrument is designed to measure the amount of stimulated electromagnetic radiation produced by pulses of electromagnetic radiation emitted into a water sample or in situ.

Dataset- specific Instrument Name	pCO2 analyzer
Generic Instrument Name	Gas Analyzer
Dataset- specific Description	pCO2 analyzer: 10-L showerhead-type seawater equilibrator (Bates et al., 1998; Sweeney et al., 2000; Takahashi et al., 1997) and an infrared CO2 detector (LI-6262, Li-Cor; Cooley et al., 2007).
Generic Instrument Description	Gas Analyzers - Instruments for determining the qualitative and quantitative composition of gas mixtures.

Dataset- specific Instrument Name	meteorological instruments on each vessel
Generic Instrument Name	Meteorological Station
Generic Instrument	MET station systems are designed to record meteorological information on board ships or mounted on moorings. These are commonly referred to as EMET (Electronic Meteorological Packages) or IMET (Improved Meteorological Packages) systems. These sensor packages record measurements of sea surface temperature and salinity, air temperature, wind speed and direction, barometric pressure, solar and long-wave radiation, humidity and precipitation.

Dataset- specific Instrument Name	
Generic Instrument Name	Niskin bottle
Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset- specific Instrument Name	Seabird thermosalinograph
Generic Instrument Name	Thermosalinograph
Generic Instrument Description	A thermosalinograph (TSG) is used to obtain a continuous record of sea surface temperature and salinity. On many research vessels the TSG is integrated into the ship's underway seawater sampling system and reported with the underway or alongtrack data.

[table of contents | back to top]

Deployments

KN197-08

Website	https://www.bco-dmo.org/deployment/58043
Platform	R/V Knorr
Report	http://bcodata.whoi.edu/ANACONDAS/ANACONDAS1-FullCruiseReport.pdf
Start Date	2010-05-22
End Date	2010-06-24
Description	ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom SymbiosesROCA: River Ocean Continuum of the Amazon WHOI cruise planning synopsis Cruise information and original data are available from the NSF R2R data catalog. Cruise Track over Salinity Climatology (Image from Dr. Ajit Subramaniam)

MV1110

Website	https://www.bco-dmo.org/deployment/58945
Platform	R/V Melville
Start Date	2011-09-03
End Date	2011-10-08
Description	ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom SymbiosesROCA: River Ocean Continuum of the Amazon Original data are available from the NSF R2R data catalog Cruise Track over Salinity Climatology (Image from Dr. Ajit Subramaniam)

AT21-04

Website	https://www.bco-dmo.org/deployment/58944
Platform	R/V Atlantis
Start Date	2012-07-13
End Date	2012-07-29
Description	ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom SymbiosesROCA: River Ocean Continuum of the Amazon WHOI cruise planning synopsis Original data are available from the NSF R2R data catalog Cruise Track over Salinity Climatology (Image from Dr. Ajit Subramaniam)

Project Information

Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses (ANACONDAS)

Website: http://amazoncontinuum.org/

Coverage: Amazon River plume; NE coast of South America; Western Tropical North Atlantic - 15N-Equator and 60W to 45W - Region surrounding the Amazon River Plume

ANACONDAS is an IMBER endorsed project. View list of all IMBER endorsed projects

View the ANACONDAS project GCMD DIF record

The ANACONDAS project was funded as part of the US National Science Foundation (NSF) Emerging Topics in Biogeochemical Cycles (ETBC) program (Directorate for Geosciences, NSF 07 -049, September 19, 2007) explicitly intended to support emerging areas of interdisciplinary research. The ETBC program aimed to foster transformational advances in the quantitative or mechanistic understanding of biogeochemical cycles that integrated physical-chemical-biological processes over the range of temporal and/or spatial scales in Earth's environments. The program especially sought proposals that addressed emerging topics in biogeochemical cycles, the water cycle or their coupling, across the interfaces of atmosphere, land, and oceans.

The ANACONDAS investigators hypothesize that large tropical river plumes with low N: P ratios provide an ideal niche for diatom-diazotroph assemblages (DDAs). They suggest that the ability of these organisms to fix N2 within the surface ocean is responsible for significant C export in the Amazon River plume. Their previous observations in the Amazon River plume helped reveal that blooms comprised of the endosymbiotic N2-fixing cyanobacterium Richelia and its diatom hosts (e.g. Hemiaulus) were a significant source of new production and carbon export. The previous work focused largely on the sensitivity of DDAs to external forcing from dust and riverine inputs, so the ecology of these organisms and the fate of their new production were largely unstudied. It is now known that DDAs are responsible for a significant amount of CO2 drawdown in the Amazon River plume, and floating sediment traps at 200 m measured 4x higher mass fluxes beneath the plume than outside the plume. This led the researchers to hypothesize that this greater export is due either to aggregation and sinking of DDAs themselves or to grazing of DDAs by zooplankton.

In this study the researchers will undertake a suite of field, satellite and modeling studies aimed at understanding the ecology and tracing the fate of C and N fixed by DDAs and other phytoplankton living in the plume. By examining C and silicate (Si) export from offshore surface waters, through the upper oceanic food web, the mesopelagic, and down to the deep sea floor, they will quantify the impact of the Amazon River on biological processes that control C sequestration and the implications of these regional processes on C, N and Si budgets. The study will go beyond previous research because they will quantify 1) the distribution, nutrient demands, and activity of DDAs in the context of phytoplankton species succession, 2) the sensitivity of the CO2 drawdown to the mix of phytoplankton, 3) the grazing and aggregation processes contributing to the sinking flux, 4) the composition of this flux, and 5) the proportion of this material that reaches the seafloor. This effort truly represents a measure of C sequestration and pump efficiency. Ecological modeling will be used to place observational results from field studies and satellites into the context of the larger Atlantic basin with tropical climate variability on interannual and longer time scales.

Three cruises were carried out during the ANACONDAS project:

AN10/KN197-08 - R/V KNORR - May/June 2010 - <u>Cruise Track over Salinity Climatology</u> (*Image: Yager, et al, 2007*)

AN11/MV1110 - R/V MELVILLE - September/October 2011 - <u>Cruise Track over Salinity Climatology</u> (*Image: Yager, et al, 2007*)

AN12/AT21-04 - R/V ATLANTIS - July/2012 - <u>Cruise Track over Salinity Climatology</u> (Image: Yager, et al, 2007)

The ANACONDAS project builds on observations made by MANTRA/PIRANA in 2001 and 2003 (RV Knorr and Seward Johnson I cruises to the same region) to address specifically 1) how carbon cycling and sequestration in the western tropical North Atlantic (WTNA) is influenced by the Amazon River through its impact on pelagic ecosystem dynamics and 2) the sensitivity of this ecosystem to anthropogenic climate change. PIRANA

revealed the importance of both riverine and atmospheric inputs for driving the high productivity of the WTNA through N2-fixation, and demonstrated the significance of the region to basin-wide biogeochemistry and C cycling. ANACONDAS will now focus on what drives phytoplankton community succession through the plume, light and nutrient requirements, factors limiting productivity, and the fate of production. These components are critical to understand the role of the plume in the regional C cycle, and to predict its response to climate variability and change.

The NSF-funded ANACONDAS project will also serve as a platform for additional measurements supported by the Gordon and Betty Moore Foundation's Marine Microbiology Initiative. ROCA (River-Ocean Continuum of the Amazon) brings additional focus on marine microbial community structure and activities, along with high-resolution measurements of organic matter along the river-ocean continuum.

ANACONDAS: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses **ROCA:** River Ocean Continuum of the Amazon

The project is funded by NSF-OCE-0934095 and NSF-OCE-0934036: Collaborative Research: ETBC: Amazon iNfluence on the Atlantic: CarbOn export from Nitrogen fixation by DiAtom Symbioses and by the Gordon and Betty Moore Foundation through GBMF-MMI-2293: River Ocean Continuum of the Amazon.

Planned Cruise Sampling

Water Column Characterization (hydrographic sampling with CTD/Rosette):

Nutrient (NO2, NO2+NO3, PO4, SiO4) concentrations

Chlorophyll a and pigments concentrations

Inorganic carbon (discrete DIC, ALK, underway pCO2)

Organic carbon, nitrogen, phosphorus

Phytoplankton and Diazotroph Abundance (using rosette and also small nets to collect)

Carbon and Nitrogen Fixation by plankton

Kinetic and Physiological Measurements of phytoplankton

Stable Isotopic Measurements of particulate material

Microbial heterotrophy

Microbial community structure and gene expression

Organic carbon and biomarker characterization

MOCNESS tows for zooplankton

Zooplankton collection for abundance and biomass Zooplankton grazing and POC flux measurements

Multicorer for deep sea sediment analyses

Solid phase analysis Pore water chemistry Isotopic composition (Pb, Th, C)

Other instrumentation over the side:

The in-water light field will be characterized with a free-falling 14 channel spectroradiometer Two "Carbon Explorers" - autonomous Sounding Oceanographic Lagrangian Observer profilers Sediment Trap Studies (using 48h deployments of floating Particle Interceptor Traps; PITs) Surface water pumps - directly bring large volumes of surface water to the deck of the ship for processing.

Shipboard Instrumentation:

ADCP 75 kHz

Bathymetry System 12 kHz

Bathymetry System 3.5 kHz

Deionized Water System

Fume Hood

HiSeas Net

Multibeam

Uncontaminated Seawater System

CTD/Water Sampling: 911+ Rosette 24-position, 10-liter bottle Rosette with dual T/C sensors

Biospherical underwater PAR (1000m depth limit)

SBE43 oxygen sensor

Wet Labs C*Star transmissometer (660nm wavelength)

Wet Labs ECO-AFL fluorometer

Dissolved Oxygen Titration System (Portable modified Winkler titration system)

Program Information

Marine Microbiology Initiative (MMI)

Website: https://www.moore.org/initiative-strategy-detail?initiativeId=marine-microbiology-initiative

A Gordon and Betty Moore Foundation Program.

Forging a new paradigm in marine microbial ecology:

Microbes in the ocean produce half of the oxygen on the planet and remove vast amounts of carbon dioxide, a greenhouse gas, from the atmosphere. Yet, we have known surprisingly little about these microscopic organisms. As we discover answers to some long-standing puzzles about the roles that marine microorganisms play in supporting the ocean's food webs and driving global elemental cycles, we realized that we still need to learn much more about what these organisms do and how they do it—including how they evolved and contribute to our ocean's health and productivity.

The Marine Microbiology Initiative seeks to gain a comprehensive understanding of marine microbial communities, including their diversity, functions and behaviors; their ecological roles; and their origins and evolution. Our focus has been to enable researchers to uncover the principles that govern the interactions among microbes and that govern microbially mediated nutrient flow in the sea. To address these opportunities, we support leaders in the field through investigator awards, multidisciplinary team research projects, and efforts to create resources of broad use to the research community. We also support development of new instrumentation, tools, technologies and genetic approaches.

Through the efforts of many scientists from around the world, the initiative has been catalyzing new science through advances in methods and technology, and to reduce interdisciplinary barriers slowing progress. With our support, researchers are quantifying nutrient pools in the ocean, deciphering the genetic and biochemical bases of microbial metabolism, and understanding how microbes interact with one another. The initiative has five grant portfolios:

Individual investigator awards for current and emerging leaders in the field.

Multidisciplinary projects that support collaboration across disciplines.

New instrumentation, tools and technology that enable scientists to ask new questions in ways previously not possible.

Community resource efforts that fund the creation and sharing of data and the development of tools, methods and infrastructure of widespread utility.

Projects that advance genetic tools to enable development of experimental model systems in marine microbial ecology.

We also bring together scientists to discuss timely subjects and to facilitate scientific exchange.

Our path to marine microbial ecology was a confluence of new technology that could accelerate science and an opportunity to support a field that was not well funded relative to potential impact. Around the time we began this work in 2004, the life sciences were entering a new era of DNA sequencing and genomics, expanding possibilities for scientific research – including the nascent field of marine microbial ecology. Through conversations with pioneers inside and outside the field, an opportunity was identified: to apply these new sequencing tools to advance knowledge of marine microbial communities and reveal how they support and influence ocean systems.

After many years of success, we will wind down this effort and close the initiative in 2021. We will have invested more than \$250 million over 17 years to deepen understanding of the diversity, ecological activities and evolution of marine microbial communities. Thanks to the work of hundreds of scientists and others involved with the initiative, the goals have been achieved and the field has been profoundly enriched; it is now positioned to address new scientific questions using innovative technologies and methods.

Integrated Marine Biogeochemistry and Ecosystem Research - US (IMBER-US)

Website: http://www.imber.info/

Coverage: global

The BCO-DMO database includes data from IMBER endorsed projects lead by US funded investigators. There is no dedicated US IMBER project or data management office. Those functions are provided by US-OCB and BCO-DMO respectively.

The information in this program description pertains to the Internationally coordinated IMBER research program. The projects contributing data to the BCO-DMO database are those funded by US NSF only. The full IMBER data catalog is hosted at the Global Change Master Directory (GCMD).

IMBER Data Portal: The IMBER project has chosen to create a metadata portal hosted by the NASA's Global Change Master Directory (GCMD). The GCMD IMBER data catalog provides an overview of all IMBER endorsed and related projects and links to datasets, and can be found at URL http://gcmd.nasa.gov/portals/imber/.

IMBER research will seek to identify the mechanisms by which marine life influences marine biogeochemical cycles, and how these, in turn, influence marine ecosystems. Central to the IMBER goal is the development of a predictive understanding of how marine biogeochemical cycles and ecosystems respond to complex forcings, such as large-scale climatic variations, changing physical dynamics, carbon cycle chemistry and nutrient fluxes, and the impacts of marine harvesting. Changes in marine biogeochemical cycles and ecosystems due to global change will also have consequences for the broader Earth System. An even greater challenge will be drawing together the natural and social science communities to study some of the key impacts and feedbacks between the marine and human systems.

To address the IMBER goal, four scientific themes, each including several issues, have been identified for the IMBER project: Theme 1 - Interactions between Biogeochemical Cycles and Marine Food Webs; Theme 2 - Sensitivity to Global Change: How will key marine biogeochemical cycles, ecosystems and their interactions, respond to global change?; Theme 3 - Feedback to the Earth System: What are the roles of the ocean biogeochemistry and ecosystems in regulating climate?; and Theme 4 - Responses of Society: What are the relationships between marine biogeochemical cycles, ecosystems, and the human system?

Ocean Carbon and Biogeochemistry (OCB)

Website: http://us-ocb.org/

Coverage: Global

The Ocean Carbon and Biogeochemistry (OCB) program focuses on the ocean's role as a component of the global Earth system, bringing together research in geochemistry, ocean physics, and ecology that inform on and advance our understanding of ocean biogeochemistry. The overall program goals are to promote, plan, and coordinate collaborative, multidisciplinary research opportunities within the U.S. research community and with international partners. Important OCB-related activities currently include: the Ocean Carbon and Climate Change (OCCC) and the North American Carbon Program (NACP); U.S. contributions to IMBER, SOLAS, CARBOOCEAN; and numerous U.S. single-investigator and medium-size research projects funded by U.S. federal agencies including NASA, NOAA, and NSF.

The scientific mission of OCB is to study the evolving role of the ocean in the global carbon cycle, in the face of environmental variability and change through studies of marine biogeochemical cycles and associated ecosystems.

The overarching OCB science themes include improved understanding and prediction of: 1) oceanic uptake and release of atmospheric CO2 and other greenhouse gases and 2) environmental sensitivities of biogeochemical cycles, marine ecosystems, and interactions between the two.

The OCB Research Priorities (updated January 2012) include: ocean acidification; terrestrial/coastal carbon

fluxes and exchanges; climate sensitivities of and change in ecosystem structure and associated impacts on biogeochemical cycles; mesopelagic ecological and biogeochemical interactions; benthic-pelagic feedbacks on biogeochemical cycles; ocean carbon uptake and storage; and expanding low-oxygen conditions in the coastal and open oceans.

Emerging Topics in Biogeochemical Cycles (ETBC)

Website: http://www.nsf.gov/pubs/2007/nsf07049/nsf07049.jsp

Coverage: global

The original call for proposals for Emerging Topics in Biogeochemical Cycles (ETBC) was issued in September 2007 by the US NSF Directorate for Geosciences (NSF 07-049).

The Geosciences Directorate (GEO) is substantially augmenting our past funding sources to explicitly support emerging areas of interdisciplinary research. We seek to foster transformational advances in our quantitative or mechanistic understanding of biogeochemical cycles that integrate physical-chemical-biological processes over the range of temporal and/or spatial scales in Earthâ \in TMs environments. We encourage submission of proposals that address emerging topics in biogeochemical cycles, the water cycle or their coupling, across the interfaces of atmosphere, land, and oceans. Proposals must cross the disciplinary boundaries of two or more divisions in Geosciences (e.g. ATM, EAR, OCE) or of at least one division in Geosciences and a division in another NSF directorate.

Although funding programmatic disciplines continues to provide a robust and adaptable framework to build our understanding of the geosciences and the earth as a system, there are classes of emerging and challenging problems that require integration of concepts and observations across diverse fields. Our goal is to enhance such integration. Successful proposals need to develop intellectual excitement in the participating disciplinary communities. Also encouraged are proposals that have broader educational, diversity, societal, or infrastructure impacts that capitalize on this interdisciplinary opportunity.

[table of contents | back to top]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0934095
Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)	<u>GBMF2293</u>
Gordon and Betty Moore Foundation: Marine Microbiology Initiative (MMI)	<u>GBMF2928</u>
NSF Division of Ocean Sciences (NSF OCE)	OCE-1133277

[table of contents | back to top]