

ZooSCAN output from of imaged zooplankton collected during BATS MOCNESS tows during R/V Atlantic Explorer cruises AE1614, AE1712, AE1830, and AE1819 in the vicinity of the Bermuda Atlantic Time-series Study from 2016 to 2018

Website: <https://www.bco-dmo.org/dataset/857891>

Data Type: Cruise Results

Version: 1

Version Date: 2021-10-05

Project

- » [Collaborative Research: Diel physiological rhythms in a tropical oceanic copepod](#) (Zooplankton Diel Rhythm)
- » [Quantifying the drivers of midwater zooplankton community structure](#) (Zooplankton Gradients)
- » [Bermuda Institute of Ocean Sciences Simons Collaboration on Ocean Processes and Ecology](#) (BIOSSCOPE)

Contributors	Affiliation	Role
Blanco-Bercial, Leocadio	Bermuda Institute of Ocean Sciences (BIOS)	Principal Investigator
Maas, Amy	Bermuda Institute of Ocean Sciences (BIOS)	Co-Principal Investigator
Gossner, Hannah	Bermuda Institute of Ocean Sciences (BIOS)	Technician
York, Amber D.	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

ZooSCAN output from of imaged zooplankton collected during BATS MOCNESS tows during R/V Atlantic Explorer cruises AE1614, AE1712, AE1830, and AE1819 in the vicinity of the Bermuda Atlantic Time-series Study from 2016 to 2018. These data were published in Maas et al. (2021).

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:32.3065 E:-64.0621 S:31.666 W:-64.7935

Temporal Extent: 2016-07-11 - 2018-10-30

Methods & Sampling

Methodology:

To obtain samples, a 1 m Multiple Opening/Closing Net and Environmental Sensing System (MOCNESS; Wiebe et al., 1985) equipped with 150 m nets was deployed during the mid-day and mid-night on cruises carried out in the vicinity of the Bermuda Atlantic Time-series Study site (BATS) in July of 2016, 2017, and 2018 as well as October 2018 (eight casts in total, 63 discrete nets; Table 1). The net sampling plan used adaptive profiling,

modifying the depth of closure to capture eight distinct ecological zones from surface to 1000 m depth: e.g. the thermocline, deep chl-a maximum (DCM), above the O₂ minimum, within the O₂ minimum core, and below the O₂ minimum (Maas et al., 2014, Steinberg et al., 2008a). The actual tow depths were set on a cruise-by-cruise basis, using CTD profiles to determine the depths of each ecological zone. Upon retrieval, the catch from each of the eight discrete nets were divided into splits. Half were preserved in 95% undenatured ethanol and the remainder preserved in buffered 4% formalin in seawater.

A representative subsample of the zooplankton community from each net were imaged from the formalin-preserved samples, then measured, using a ZooSCAN ver. 3 at 4,800 dpi (following the methods in: Gorsky et al., 2010, Vandromme et al., 2012). In order to better represent all size classes in the images, the original sample was divided into three size categories. All individuals larger than 2 cm were selected by eye and scanned separately from all the others. The remainder of the sample was sieved through a 1-mm mesh sieve, and both size fractions were individually scanned. From these smaller size fractions, at least 1500 particles were scanned after subsampling using a Motoda splitter (Motoda, 1959), requiring generation of two separate scans for both size classes. This resulted in a total of five images per net. Raw images were then processed in ZooProcess (Gorsky et al., 2010, Vandromme et al., 2012), and the resulting vignettes, along with a measurement and metadata file, were uploaded to EcoTaxa (<https://ecotaxa.obs-vlfr.fr/>; Picheral et al., 2017) for machine-assisted identification, using a training set developed by the authors. The taxonomy of one paired set of tows was completely validated (July 2017). For the remaining tows taxonomy is based on the Ecotaxa prediction using this 2017 training set.

Object_id and imagenames:

Related Datasets may contain the image name which is constructed the same way as the object_id in this dataset except it does not have the additional _# at the end, and the imagename ends in the .tif extension. This additional number in the object_id is added by the Zooprocess software (Hydroptic, 2016).

e.g.

object_id: ae1614_m3_n1_d2_a_1_100

image_name: ae1614_m3_n1_d2_a_1.tif

Instruments: ZooSCAN ver. 3, 1m MOCNESS

Problem report: During the July 2018 cruise the top net of the night cast failed, and consequently data from this net is absent.

Data Processing Description

BCO-DMO data manager processing notes:

- * File DATASET_M3_to_M13_BATS_v3Nov21.txt imported into the BCO-DMO data system.
- * zero padding time for cases when single digit hour (e.g. 35000-> 035000 meaning 03:50:00).
- * local dates and time converted to ISO 8601 formats (yyyy-mm-dd and HH:MM:SS)
- * Added timestamp in ISO 8601 format yyyy-mm-ddTHH:MM:SSZ

[[table of contents](#) | [back to top](#)]

Data Files

File
zooscan_mocness_output.csv (Comma Separated Values (.csv), 31.51 MB) MD5:5205c187d64857fe84cb67839ad2f6be
Primary data file for dataset ID 857891

[[table of contents](#) | [back to top](#)]

Supplemental Files

File

Image Sample Metadata (ZooSCAN BATS M3 to M13)

filename: image_sample_information.csv

(Comma Separated Values (.csv), 30.56 KB)
MD5:1954f230c158230d25e019f37d82c19a

Image Sample Metadata for ZooSCAN images collected during R/V Atlantic Explorer cruises AE1614, AE1712, AE1819, AE1830 between 2016-2018.

Parameters (column name, description, units):

image_name,Image name, unitless

cruise_id,Cruise identifier, unitless

mocness_id,MOCNESS identifier,unitless

net_id,Net identifier,unitless

size_fraction,Size fraction (e.g. d3). See methodology,unitless

replicate,Replicate (see methodology),unitless

lat,Latitude,decimal degrees

lon,Longitude,decimal degrees

date, Date (local) in time zone AST (UTC-4)/ADT (UTC-3) in format YYYY-MM-DD,unitless

time, Time (local) in time zone AST (UTC-4)/ADT (UTC-3) in format hh:mm:ss

ISO_DateTime_UTC,Datetime in timezone UTC in ISO 8601 format YYYY-MM-DDThh:mm:ssZ

[[table of contents](#) | [back to top](#)]

Related Publications

Gorsky, G., Ohman, M. D., Picheral, M., Gasparini, S., Stemmann, L., Romagnan, J.-B., ... Prejger, F. (2010). Digital zooplankton image analysis using the ZooScan integrated system. *Journal of Plankton Research*, 32(3), 285–303. doi:[10.1093/plankt/fbp124](https://doi.org/10.1093/plankt/fbp124)

Methods

Hydroptic (2016). ZooSCAN. Available at

http://www.hydroptic.com/index.php/public/Page/product_item/ZOOSCAN. Accessed June 17th, 2021.

Software

Maas, A. E., Frazar, S. L., Outram, D. M., Seibel, B. A., & Wishner, K. F. (2014). Fine-scale vertical distribution of macroplankton and micronekton in the Eastern Tropical North Pacific in association with an oxygen minimum zone. *Journal of Plankton Research*, 36(6), 1557–1575. doi:[10.1093/plankt/fbu077](https://doi.org/10.1093/plankt/fbu077)

Methods

Maas, A. E., Gossner, H., Smith, M. J., & Blanco-Bercial, L. (2021). Use of optical imaging datasets to assess biogeochemical contributions of the mesozooplankton. *Journal of Plankton Research*, 43(3), 475–491.

doi:[10.1093/plankt/fbab037](https://doi.org/10.1093/plankt/fbab037)

Results

Motoda, S. (1959) Devices of simple plankton apparatus. *Memoirs of the Faculty of Fisheries Hokkaido University*, 7, 73-94. Available from <http://hdl.handle.net/2115/21829>.

Methods

Picheral M, Colin S, Irisson J-O (2017). EcoTaxa, a tool for the taxonomic classification of images.

<http://ecotaxa.obs-vlfr.fr>

IsDerivedFrom

Steinberg, D. K., Cope, J. S., Wilson, S. E., & Kobari, T. (2008). A comparison of mesopelagic mesozooplankton community structure in the subtropical and subarctic North Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 55(14-15), 1615–1635. doi:[10.1016/j.dsr2.2008.04.025](https://doi.org/10.1016/j.dsr2.2008.04.025)

Methods

Vandromme, P., Stemmann, L., Garcia-Comas, C., Berline, L., Sun, X., & Gorsky, G. (2012). Assessing biases in computing size spectra of automatically classified zooplankton from imaging systems: A case study with the

ZooScan integrated system. *Methods in Oceanography*, 1-2, 3-21. doi:[10.1016/j.mio.2012.06.001](https://doi.org/10.1016/j.mio.2012.06.001)
Methods

Wiebe, P. H., Morton, A. W., Bradley, A. M., Backus, R. H., Craddock, J. E., Barber, V., ... Flierl, G. R. (1985). New development in the MOCNESS, an apparatus for sampling zooplankton and micronekton. *Marine Biology*, 87(3), 313-323. doi:10.1007/bf00397811 <https://doi.org/10.1007/BF00397811>
Methods

[[table of contents](#) | [back to top](#)]

Related Datasets

References

Blanco-Bercial, L., Maas, A., Gossner, H. (2021) **ZooSCAN biovolume to biomass from imaged zooplankton collected during MOCNESS tows during various R/V Atlantic Explorer cruises and small boat deployments in the Sargasso Sea between 2016 to 2019**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-06-17
doi:10.26008/1912/bco-dmo.854077.1 [[view at BCO-DMO](#)]
Relationship Description: Biovolume:biomass conversion provides validation for the analysis of the ZooScan output.

IsRelatedTo

Blanco-Bercial, L., Maas, A., Gossner, H. (2021) **ZooSCAN images of zooplankton collected during BATS MOCNESS tows during R/V Atlantic Explorer cruises AE1614, AE1712, AE1830, and AE1819 in the vicinity of the Bermuda Atlantic Time-series Study from 2016 to 2018**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2021-10-07
doi:10.26008/1912/bco-dmo.853440.1 [[view at BCO-DMO](#)]
Relationship Description: Images for the ZooScan output in this dataset.

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
object_lat	Latitude of collection	decimal degrees
object_lon	Longitude of collection	decimal degrees
object_date	Date of Collection in format yyyyymmdd	unitless
object_time	Time of Collection in format hhmmss	unitless
object_depth_min	Minimum depth of net collection	m
object_depth_max	Maximum depth of net collection	m
object_annotation_hierarchy	Taxonomic classification (predicted except for AE1712 which was validated)	unitless
object_area	Object area in pixels	pix ²
object_major	Object major axis length in pixels	pix
object_minor	Object minor axis length in pixels	pix
object_feret	Object Maximum feret diameter, i.e., the longest distance between any two points along the object boundary in pixels	pix
object_esd	Object Equivalent Spherical Diameter	pix
sample_tot_vol	Volume filtered	m ³
acq_sub_part	number of splits	number of splits
object_id	Particle identifier (cruise#_mocnessID_net#_sizefraction_image#)	
ISO_DateTime_UTC	Datetime (UTC) of Collection in ISO 8601 format yyyy-mm-ddTHH:MMZ.	%Y-%m-%dT%H:%M:%SZ

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	1m MOCNESS
Generic Instrument Name	MOCNESS
Generic Instrument Description	The Multiple Opening/Closing Net and Environmental Sensing System or MOCNESS is a family of net systems based on the Tucker Trawl principle. There are currently 8 different sizes of MOCNESS in existence which are designed for capture of different size ranges of zooplankton and micro-nekton Each system is designated according to the size of the net mouth opening and in two cases, the number of nets it carries. The original MOCNESS (Wiebe et al, 1976) was a redesigned and improved version of a system described by Frost and McCrone (1974).(from MOCNESS manual) This designation is used when the specific type of MOCNESS (number and size of nets) was not specified by the contributing investigator.

Dataset-specific Instrument Name	ZooSCAN ver. 3
Generic Instrument Name	ZooSCAN
Dataset-specific Description	http://www.hydroptic.com/index.php/public/Page/product_item/ZOOSCAN
Generic Instrument Description	Description excerpt from Hydroptic website http://www.hydroptic.com/index.php/public/Page/product_item/ZOOSCAN The ZooSCAN (CNRS patent) system makes use of scanner technology with custom lighting and a watertight scanning chamber into which liquid zooplankton samples can be placed. The scanner recovers a high-resolution, digital image and the sample can be recovered without damage. These digital images can then be investigated by computer processing. While the resolution of the digitized zooplankton images is lower than the image obtained using a binocular microscope this technique has proved to be more than adequate for large sample sets. Identification of species is done by automatic comparison of the image (vignette) of each individual animal in the scanned image with a library data set which may be built by the investigator for each individual survey or imported from a previous survey. The latest machine learning algorithm allows high recognition levels even if we recommend complementary manual sorting to achieve a high number of taxonomic groups.

[[table of contents](#) | [back to top](#)]

Deployments

AE1614

Website	https://www.bco-dmo.org/deployment/853444
Platform	R/V Atlantic Explorer
Report	https://datadocs.bco-dmo.org/docs/305/BIOSCOPE/data_docs/AE1614_CS_narrative_FINAL.pdf
Start Date	2016-07-09
End Date	2016-07-12

AE1712

Website	https://www.bco-dmo.org/deployment/857372
Platform	R/V Atlantic Explorer
Report	https://datadocs.bco-dmo.org/docs/305/BIOSCOPE/data_docs/AE1712_CS_narrative.pdf
Start Date	2017-07-08
End Date	2017-07-11
Description	Project BIOS-SCOPE

AE1830

Website	https://www.bco-dmo.org/deployment/857780
Platform	R/V Atlantic Explorer
Start Date	2018-10-27
End Date	2018-10-31
Description	Cruise DOI: 10.7284/908188

AE1819

Website	https://www.bco-dmo.org/deployment/857784
Platform	R/V Atlantic Explorer
Report	https://datadocs.bco-dmo.org/docs/305/BIOSCOPE/data_docs/AE1819_CS_narrative_v1.pdf
Start Date	2018-07-03
End Date	2018-07-06
Description	Project BIOS-SCOPE

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Diel physiological rhythms in a tropical oceanic copepod (Zooplankton Diel Rhythm)

Coverage: Bermuda

NSF Award Abstract:

The daily vertical migration (DMV) of zooplankton and fish across hundreds of meters between shallow and deep waters is a predominant pattern in pelagic ecosystems. This migration has consequences for biogeochemical cycling as it moves a substantial portion of fixed carbon and nitrogen (an estimated 15 to 40 % of the total global organic export) from the surface directly to depth where it feeds the midwater food chain and sequesters nutrients away from atmospheric mixing. Estimates and predictions of these fluxes are, however, poorly understood at present. New observations have shown that one source of uncertainty is due to the assumption that metabolic rates and processes do not vary over the course of the day, except based on changes in temperature and oxygen availability. Rates are, however, also driven by differences in feeding, swimming behavior, and underlying circadian cycles. The objective of this project is to improve the ability of scientists to understand and predict zooplankton contributions to the movement of carbon and nitrogen in the ocean by detailing daily changes in physiological processes of these organisms. By producing a set of respiration and excretion measurements over a daily time series, paired with simultaneously collected gene and protein expression patterns for an abundant vertically migratory species, the investigators will provide unprecedented and predictive insight into how changes in the environment affect the contribution of zooplankton to biogeochemical fluxes. The sampling design of the project will advance discovery and understanding by providing hands-on training opportunities to at least two undergraduate researchers. The project will broaden dissemination of the research via development of an educational module, focusing on rhythms in the ocean. The module will initially be piloted with the Bermuda Institute of Ocean Sciences (BIOS) summer camp students and then disseminated through the BIOS Explorer program, the Teacher Resources Page on the BIOS website, and published in a peer-reviewed educational journal.

This project will characterize the metabolic consequences of daily physiological rhythms and DVM for a model zooplankton species, the abundant subtropical copepod *Pleuromamma xiphioides*. Flux processes (oxygen consumption, carbon dioxide production, production of ammonium and fecal pellet production) will be interrogated using directed experiments testing the effects of temperature, feeding and circadian cycle. Circadian cycling will further be examined using transcriptomic and proteomic profiling. These experiments will be related to field samples taken at 6-h intervals over the course of the diel migration using an integrated suite

of molecular and organismal metrics. Combined organismal, transcriptomic and proteomic profiles will provide an understanding of which metabolic pathways and associated flux products vary in relation to particular environmental variables (food, light cycle, temperature). Diel variation in metabolic rates will also be assessed across seasons and species using other important migratory groups (pteropod, euphausiid, and another copepod). The metabolic data will then be contextualized with abundance estimates from archived depth-stratified tows to allow scaling to community-level patterns and will be used to improve calculations of zooplankton contribution to particulate organic carbon, nitrogen and respiratory active flux. The results of this study will both improve our flux estimates and provide predictive insight into how various environmental variables influence the underlying physiological pathways generating carbon and nitrogen flux.

Cruise reports are available from the completed cruises:

[SD031019](#)

[AE1910](#)

[AE1918](#)

Quantifying the drivers of midwater zooplankton community structure (Zooplankton Gradients)

Website: https://www.nsf.gov/awardsearch/showAward?AWD_ID=1948162

Coverage: North Atlantic and North Pacific

NSF Award Abstract:

Processes in the midwater region below 200 m depth, also known as the twilight zone, represent a major unknown for the biology and chemistry of the ocean. Studies of animals drifting in the oceans, known as zooplankton, are scarce due to the difficulty and associated time and costs of sampling deep waters. The advent of automated image analysis and genetic tools is leading to a rapid increase in our knowledge of the diversity, abundances and size distribution of communities in shallow waters. However, our understanding of the deeper layers of the ocean is still in its infancy, and there are few studies that combine these three facets of the ecology of the zooplankton. The objective of this project is to leverage existing samples, obtained from previously NSF-funded research in the North Pacific and North Atlantic, to study how the abundances, diversity, and size distribution of zooplankton in the midwater vary with latitude and environmental factors. Automated image analyses provide information on abundance and size, and genetic analyses give unprecedented data on the diversity of the midwater community for the North Atlantic and the North Pacific, from subtropical to subarctic environments. This project provides high quality hands-on training opportunities for at least two undergraduate researchers and generates material for undergraduate and graduate courses. Two workshops train educators on the classroom use of the NSF-funded Biological and Chemical Oceanography Data Management Office (BCO-DMO) open access oceanographic data.

Recent advances in image analysis and metabarcoding of zooplankton communities via new data tools are an opportunity to generate quantitative and predictive relationships between environmental drivers and zooplankton diversity, abundances and size distribution. While this information is available for plankton in epipelagic regions, the focus of this study is on midwater communities, which remain poorly characterized. Obtaining these data is the first step towards a quantitative analysis that assesses the impact of the midwater community on biogeochemical cycles. The project uses archived samples from two cruises conducted in the N. Atlantic and N. Pacific to test hypotheses about how temperature, midwater hypoxia, primary productivity and biogeographic province shape the size class structure, biodiversity and behavior (diurnal vertical migration) of zooplankton communities. These newly-generated image and metabarcoding datasets of the mesozooplankton community from 0-1000 m are cross-comparable with other ocean regions. These data describe how migratory and midwater resident zooplankton communities are structured by environmental variables and demonstrate how this influences their biogeochemical contributions (specifically active flux and midwater attenuation of flux). Data tools generated for the image analysis in combination with metabarcoding has broad application in plankton ecology and allows metaanalysis of other datasets. The project is complementary to ongoing national and international projects that seek to describe the function and structure of the midwater. In contrast to existing modeling and process projects, this project covers a moderately large geographic area and thus provides a strong comparative foundation for broader community-wide assessment of the function of zooplankton in the twilight zone.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using

the Foundation's intellectual merit and broader impacts review criteria.

Bermuda Institute of Ocean Sciences Simons Collaboration on Ocean Processes and Ecology (BIOSSCOPE)

Website: <http://scope.bios.edu/>

Coverage: North Atlantic Subtropical Gyre, Bermuda Atlantic Time Series (BATS) site

The aim of BIOS-SCOPE is to expand knowledge about the BATS ecosystem and achieve a better understanding of ocean food web sources, sinks and transformations of DOM. Advances in knowledge and technology now pose us to investigate the specific mechanisms of DOM incorporation, oxidation and transformation by zooplankton and the distinct microbial plankton communities that have been discovered at BATS.

The overarching goal of the BIOS-SCOPE is to form and foster collaborations of cross disciplinary science that utilize a broad suite of genomic, chemical, ecological, and biogeochemical approaches to evaluate microbial process, structure and function on various scales. These scales will range from organism-compound and organism-organism interactions to large biogeochemical patterns on the ecosystem scale. For this purpose we have assembled a cross-disciplinary team including microbial oceanographers (Carlson and Giovannoni), a chemical oceanographer (Kujawinski), biological oceanographer / zooplankton ecologists (Maas and Blanco-Bercial) and microbial bioinformatician (Temperton) with the expertise and technical acuity that are needed to study complex interactions between food web processes, microbes and DOM quantity and quality in the oligotrophic ocean. This scientific team has a vision of harnessing this potential to produce new discoveries that provide a mechanistic understanding of the carbon cycle and explain the many emergent phenomenon that have yet to be understood.

For additional details:

- **BIOS-SCOPE Narrative:**
https://datadocs.bco-dmo.org/docs/302/BIOSSCOPE/data_docs/BIOS-SCOPE_Narrative_FINAL.pdf
- **Physical Framework:**
https://datadocs.bco-dmo.org/docs/302/BIOSSCOPE/data_docs/Physical_Framework.pdf

BIOSSCOPE I: November 1st, 2015 through October 31st, 2020

Current: November 1st, 2020 to October 31st, 2025

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
Simons Foundation (Simons)	unknown SCOPE Simons
NSF Division of Ocean Sciences (NSF OCE)	OCE-1829318
NSF Division of Ocean Sciences (NSF OCE)	OCE-1948162

[[table of contents](#) | [back to top](#)]