Pyrosome (Pyrosoma atlanticum) counts from vertical video profiles in the Northern California Current acquired between February and July 2018.

Website: https://www.bco-dmo.org/dataset/866614 Data Type: Cruise Results Version: 1 Version Date: 2021-12-10

Project

» Collaborative Research: Mesozooplankton food webs in intermittent upwelling systems: An overlooked link in a productive ocean (MEZCAL)

Contributors	Affiliation	Role
Sutherland, Kelly Rakow	University of Oregon	Principal Investigator
Cowen, Robert K.	Oregon State University (OSU-HMSC)	Co-Principal Investigator
<u>Sponaugle, Su</u>	Oregon State University (OSU)	Co-Principal Investigator
<u>Lyle, Joanna</u>	University of Oregon (OIMB)	Contact
<u>Soenen, Karen</u>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Pyrosome (Pyrosoma atlanticum) counts from vertical video profiles in the Northern California Current acquired between February and July 2018.

Table of Contents

- <u>Coverage</u>
- Dataset Description
 - Methods & Sampling
 - Data Processing Description
- Data Files
- <u>Related Publications</u>
- <u>Parameters</u>
- Instruments
- Deployments
- Project Information
- Funding

Coverage

Spatial Extent: N:44.6532 **E**:-124.117 **S**:41.0575 **W**:-125.118 **Temporal Extent**: 2018-02-17 - 2018-07-12

Dataset Description

These data were published in Lyle, J.T. et al. 2021

Methods & Sampling

We mounted a GoPro Hero 4 (4K, 30fps) in a deep-water housing (GoDeep Aluminum, Sexton Inc.) and two 7500 lumen lights (BigBlue VL7500P) to the ship's onboard CTD rosette frame. At each station, simultaneous CTD and camera deployments captured fine-scale (1 m), *in situ* counts of pyrosomes to 100 m, or 5 m above the bottom at shallower stations. Camera frames were subsequently analyzed to determine pyrosome

distributions with depth. For each meter of depth, we extracted a still frame from the camera and counted all pyrosome colonies. We adjusted this count by subtracting colonies that were visible in the still frame of the previous meter to avoid double-counts.

In situ videos were captured using a GoPro Hero 4 (4K, 30fps) in a deep-water housing (GoDeep Aluminum, Sexton Inc.) with two 7500 lumen lights (BigBlue VL7500P) mounted to the ship's onboard CTD rosette frame. Video was aligned to depth information captured by the CTD (SBE 911).

Data Processing Description

Videos were processed using VirtualDub. Depth information was extracted from CTD files using SBE data processing (v.7.26.6).

[table of contents | back to top]

Data Files

File
pyrosomes_cam.csv(Comma Separated Values (.csv), 236.88 KB) MD5:92992337ac78592a3405b84914a78526
Primary data file for dataset ID 866614

[table of contents | back to top]

Related Publications

Lyle, J. T., Cowen, R. K., Sponaugle, S., & Sutherland, K. R. (2022). Fine-scale vertical distribution and diel migrations of Pyrosoma atlanticum in the northern California Current. Journal of Plankton Research, 44(2), 288–302. https://doi.org/<u>10.1093/plankt/fbac006</u> *Results*

[table of contents | back to top]

Parameters

Parameter	Parameter Description	
Cruise	Cruise identification number: S=summer; W=winter; 18=2018	
Location	on Sampling location: NH=Newport, OR; TR=Trinidad Head, CA	
Station	Location along transect: 1=closest nearshore; 5=furthest offshore; D=day; N=night	
Transect	Sampling transect: Ma=first transect; Mb=second transect	
Date_Time_UTC	Date and time of the start of the net towing in UTC time zone	unitless
Latitude	Latitude of deployment start location, north is positive	decimal degrees
Longitude	ongitude Longitude of deployment start location, west is negative	
Depth	Depth of sampling in meters. Depth bins start at the surface to 100 m.	
Pyro_binned_count Pyrosome counts from vertical video profiles in 1-m depth bins. Data ma missing near the surface due to visual disturbances, or at shallower stat where the bottom depth is < 100 m.		counts

Instruments

Dataset- specific Instrument Name	CTD (SBE 911)	
Generic Instrument Name	CTD Sea-Bird 911	
Dataset- specific Description	In situ videos were captured using a GoPro Hero 4 (4K, 30fps) in a deep-water housing (GoDeep Aluminum, Sexton Inc.) with two 7500 lumen lights (BigBlue VL7500P) mounted to the ship's onboard CTD rosette frame. Video was aligned to depth information captured by the CTD (SBE 911).	
Generic Instrument Description		
Dataset- specific Instrument Name	GoPro Hero 4 (4K, 30fps)	
Generic Instrument Name	Underwater Camera	
Dataset- specific Description	In situ videos were captured using a GoPro Hero 4 (4K, 30fps) in a deep-water housing (GoDeep Aluminum, Sexton Inc.) with two 7500 lumen lights (BigBlue VL7500P) mounted to the ship's onboard CTD rosette frame. Video was aligned to depth information captured by the CTD (SBE 911).	
Generic Instrument Description	All types of photographic equipment that may be deployed underwater including stills, video, film and digital systems.	

[table of contents | back to top]

Deployments

SKQ201804S		
Website	https://www.bco-dmo.org/deployment/783051	
Platform	R/V Sikuliaq	
Start Date	2018-02-17	
End Date	2018-02-23	

Website	https://www.bco-dmo.org/deployment/783078	
Platform	R/V Sally Ride	
Start Date	2018-07-06	
End Date	2018-07-11	

[table of contents | back to top]

Project Information

Collaborative Research: Mesozooplankton food webs in intermittent upwelling systems: An overlooked link in a productive ocean (MEZCAL)

Coverage: Northern California Current

This project will examine the coastal ocean mesozooplankton community and their predation by early life stages of fish in the northern California Current. The goal is to understand how these predator-prey interactions change during different oceanographic regimes that vary seasonally in the region. This study will use a very high-resolution imaging system coupled with net samples to measure trophic interactions within the zooplankton community across a range of environmental parameters (e.g., temperature, relative timing and intensity of upwelling). The camera provides detailed information on the fine-scale abundance and spatial distributions of a wide diversity of plankton, while the net samples will provide biological samples for diet-related analyses. This project will train 12 undergraduate and two graduate students and one post-doctoral scholar. The research team will develop a variety of educational activities and products to facilitate greater outreach to public audiences. Plankton imagery from this project will be used to build the Global Plankton Imagery Library, an open-access repository for plankton imagery that will be a resource for the research community. The researchers will expand the imagery available in the Plankton Portal, a public website they developed in partnership with the Citizen Science Alliance's Zooniverse, that invites citizen scientists to participate in classifying plankton from field photographs. They will collaborate with Science Education specialists to include Plankton Portal kiosks in a new public exhibit at the Oregon State University's Hatfield Marine Science Center (HMSC) Visitor Center, which annually hosts 150,000 visitors of all ages. Importantly, this activity will not only educate K-12 and beyond, but will enable researchers to study what factors motivate citizen scientists, what characterizes "heavy-users", and how those users can be supported and encouraged into advanced collaborator roles. A multi-media artist will join the research cruises as part of the new Artist-At-Sea program. Their artwork will be displayed at the HMSC Visitor Center and University of Oregon's Charleston Marine Life Center and a scaled traveling show will tour Oregon metropolitan areas and underserved communities.

Eastern boundary currents are among the most productive marine ecosystems on the planet and support a significant proportion of global fisheries, yet there are unanswered guestions about the role of non-crustacean zooplankton in transferring production through upwelling food webs. This study will answer key questions about the food web dynamics associated with mesozooplankton linkages, sources of carbon production, and consequences for upper trophic levels in different shelf upwelling systems. Not only is there a knowledge gap in how the food web currently functions in transition areas of major eastern boundary current systems, but there is increasing evidence that these systems are changing. Regional and global shifts in major currents, including upwelling strength, together with temperature-induced latitudinal shifts in species ranges that are already occurring and predicted to continue will have major effects on interactions among species, and consequently, food webs. Understanding these interactions and predicting future changes is highly relevant to science, society, and economies. The researchers plan to sample the winter and summer seasons in the northern California Current off central Oregon (intermittent upwelling) and northern California (continuous upwelling) with the high resolution In Situ Ichthyoplankton Imaging System to obtain an accurate description of mesozooplankton communities: their abundances, and horizontal and vertical spatial distributions, over contrasting upwelling/downwelling system dynamics. In parallel, they plan to collect depth-discrete mesozooplankton samples to quantify seasonal diets for larval fishes and gelatinous zooplankton and preyspecific growth rates of larval fishes. Stable isotope analysis of mesozooplankton predators and prey will reveal the relative role of new vs. regenerated production in sustaining food webs such major eastern boundary currents.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1737399</u>
NSF Division of Ocean Sciences (NSF OCE)	<u>OCE-1737364</u>

[table of contents | back to top]