

# Oxygen to argon molar ratios measured in surface waters of Station ALOHA between June 16 and June 23, 2019 onboard R/V Kilo Moana cruise KM1910.

**Website:** <https://www.bco-dmo.org/dataset/868774>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2022-05-25

## Project

» [EAGER Collaborative Research: Early career chief scientist training for biological and chemical oceanographers](#)  
(Chief Sci KM1910)

Contributors	Affiliation	Role
<a href="#">Ferrón, Sara</a>	University of Hawaii at Manoa (SOEST)	Principal Investigator, Contact
<a href="#">Soenen, Karen</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Oxygen to argon molar ratios measured in surface waters of Station ALOHA between June 16 and June 23, 2019 onboard R/V Kilo Moana cruise KM1910.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Supplemental Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Deployments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:22.83 E:-157.924 S:22.667 W:-158.045

**Temporal Extent:** 2019-06-16 - 2019-06-23

## Methods & Sampling

Samples for O<sub>2</sub>/Ar analyses were collected at three different depths within the mixed layer (5, 15, and 25 m) with 12-L Niskin® bottles attached to a CTD rosette. Seawater samples, collected in triplicate, were transferred to 12mL Labco Exetainer® screw cap vials with rubber septa. The vials were filled from the bottom using Tygon® tubing and allowed to overflow at least 3 times their volume. Samples were poisoned with 50 µL of saturated mercuric chloride solution and stored in the dark immersed in water at 4°C until analysis on board, within 2 days of collection.

## Data Processing Description

Dissolved O<sub>2</sub>/Ar was measured in discrete water samples using membrane inlet mass spectrometry as described by Ferrón et al. (2015).

Membrane inlet mass spectrometer consists of a Pfeiffer Vacuum HiCube 80 Eco turbo pumping station connected to a HiQuay™ quadrupole mass spectrometer (QMG700), with a Balzers radio frequency generator (QMH 400-5) and a Balzers analyzer (QMA 430). The membrane inlet design is from Bay Instruments (Easton, Maryland).

BCO-DMO Processing Notes:

\* Adjusted column names to comply with database requirements

\* Added ISO8601 format of date and times

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>mixed_layer.csv</b> (Comma Separated Values (.csv), 8.51 KB) MD5:1f7f5a094228e4769c491d6f78bd1812 Primary data file for dataset ID 868774

[ [table of contents](#) | [back to top](#) ]

---

## Supplemental Files

File
<b>Mixed_layer_oxygen_to_argon_km1910</b> filename: Mixed_layer_oxygen_to_argon_km1910.xlsx (Octet Stream, 18.24 KB) MD5:f5612c8f0ae962eb1ac3e13daa5ff9f8 Mixed layer oxygen to argon km1910: Data table with mean and standard deviation of three replicates of parameters Oxygen to Argon molar ratio and Biological O <sub>2</sub> saturation anomaly.

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

De Boyer Montégut, C. (2004). Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *Journal of Geophysical Research*, 109(C12). doi:10.1029/2004jc002378

<https://doi.org/doi:10.1029/2004JC002378>

*Methods*

Ferrón, S., Wilson, S. T., Martínez-García, S., Quay, P. D., & Karl, D. M. (2015). Metabolic balance in the mixed layer of the oligotrophic North Pacific Ocean from diel changes in O<sub>2</sub> /Ar saturation ratios. *Geophysical Research Letters*, 42(9), 3421–3430. doi:10.1002/2015gl063555 <https://doi.org/10.1002/2015GL063555>

*Methods*

García, H. E., & Gordon, L. I. (1992). Oxygen solubility in seawater: Better fitting equations. *Limnology and Oceanography*, 37(6), 1307–1312. doi:[10.4319/lo.1992.37.6.1307](https://doi.org/10.4319/lo.1992.37.6.1307)

*Methods*

Hamme, R. C., & Emerson, S. R. (2004). The solubility of neon, nitrogen and argon in distilled water and seawater. *Deep Sea Research Part I: Oceanographic Research Papers*, 51(11), 1517–1528.

doi:[10.1016/j.dsr.2004.06.009](https://doi.org/10.1016/j.dsr.2004.06.009)

*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Cruise_ID	Cruise identification number	unitless
Sample_ID	Sample identification number	unitless
Date	Date and time of CTD cast	UTC
Latitude	Latitude, south is negative	decimal degrees
Longitude	Longitude, west is negative	decimal degrees
Cast	Cast number	unitless
Niskin	Niskin bottle	unitless
Target_depth	Target depth	meters (m)
MLD	Mixed layer depth, calculated following de Boyer Montégut et al. (2004), as the first depth where the potential density is at least 0.03 kg m <sup>-3</sup> larger than the value at 10 m	meters (m)
Temp	Conservative temperature	degrees Celsius (°C)
Sal	Absolute salinity	g/kg
Sig	Potential density	kg/m <sup>3</sup>
O2_sol	Oxygen solubility determined from Garcia and Gordon (1992) solubility equations	mmol/m <sup>3</sup>
O2_Ar	Oxygen to Argon molar ratio.	mol O <sub>2</sub> /mol Ar
D_O2_Ar	Biological O <sub>2</sub> saturation anomaly in percent. $\delta(O_2/Ar) = [(O_2/Ar)_{measured} / (O_2/Ar)_{equilibrium} - 1] * 100$ . Oxygen solubility determined from Garcia and Gordon (1992). Argon solubility determined from Hamme and Emerson (2004)	% (percent)
ISO_DateTime_UTC	Sampling start date and time (UTC) in ISO8601 format: YYYY-MM-DDThh:mm:ssz	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Membrane inlet mass spectrometer
<b>Generic Instrument Name</b>	Membrane Inlet Mass Spectrometer
<b>Dataset-specific Description</b>	Membrane inlet mass spectrometer consists of a Pfeiffer Vacuum HiCube 80 Eco turbo pumping station connected to a HiQuay™ quadrupole mass spectrometer (QMG700), with a Balzers radio frequency generator (QMH 400-5) and a Balzers analyzer (QMA 430). The membrane inlet design is from Bay Instruments (Easton, Maryland).
<b>Generic Instrument Description</b>	Membrane-introduction mass spectrometry (MIMS) is a method of introducing analytes into the mass spectrometer's vacuum chamber via a semipermeable membrane.

## Deployments

### KM1910

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/841636">https://www.bco-dmo.org/deployment/841636</a>
<b>Platform</b>	R/V Kilo Moana
<b>Report</b>	<a href="https://datadocs.bco-dmo.org/docs/305/Chief_Sci_KM1910/data_docs/matt_church_EAGER_cruise_plan_06_17_2019.pdf">https://datadocs.bco-dmo.org/docs/305/Chief_Sci_KM1910/data_docs/matt_church_EAGER_cruise_plan_06_17_2019.pdf</a>
<b>Start Date</b>	2019-06-15
<b>End Date</b>	2019-06-24
<b>Description</b>	NSF Chief Scientist Training Cruise. For more information, see Rolling Deck to Repository (R2R): <a href="https://www.rvdata.us/search/cruise/KM1910">https://www.rvdata.us/search/cruise/KM1910</a> (cruise DOI: 10.7284/908380)

## Project Information

### **EAGER Collaborative Research: Early career chief scientist training for biological and chemical oceanographers (Chief Sci KM1910)**

**Coverage:** Station ALOHA (22.75N, 158W), North Pacific Ocean

#### *NSF Award Abstract:*

##### Intellectual Merit

The PIs request funds to provide training in leading and organizing research cruises to early career researchers in the areas of Biological and Chemical Oceanography. Participants in this training program would be introduced to pre-cruise planning and logistics, receive training in commonly used oceanographic sampling equipment, and conduct shipboard measurements during a 10-day oceanographic cruise to the North Pacific Subtropical Gyre (NPSG). The goal of this training program is to prepare early career scientists for leading and participating in interdisciplinary oceanographic research at sea.

##### Broader Impacts

The proposed program addresses the broader impacts criteria successfully. The research cruise and follow-up reports and publications focus on interdisciplinary questions important for advancing the field. Given the rapid changes that oceanic systems are undergoing, it is important to have a cadre of junior scientists who are adept at managing interdisciplinary collaborations and conducting research at sea. The PIs are considering ways to connect with diverse audiences in recruiting participants. The impact on early career oceanographers will be very strong. This will create an experience that will be a major impact on the careers of the trainees, especially if they stay in the oceanography field.

## Funding

<b>Funding Source</b>	<b>Award</b>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1911990</a>