

# Abundance of aggregates of the marine diatom *Minutocellus polymorphus* and particle-associated marine bacteria from culture and roller tank experiments in 2021-2022

**Website:** <https://www.bco-dmo.org/dataset/876461>

**Data Type:** experimental

**Version:** 1

**Version Date:** 2022-06-27

## Project

» [Aggregation of Marine Picoplankton](#) (Marine Plankton Aggregation)

Contributors	Affiliation	Role
<a href="#">Neuer, Susanne</a>	Arizona State University (ASU)	Principal Investigator
<a href="#">Cadillo-Quiroz, Hinsby</a>	Arizona State University (ASU)	Co-Principal Investigator
<a href="#">Cruz, Bianca N.</a>	Arizona State University (ASU)	Co-Principal Investigator
<a href="#">Gerlach, Dana Stuart</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Laboratory data on the formation of aggregates of the marine nanodiatom *Minutocellus polymorphus* with and without the presence of known particle-associated and aggregation-enhancing marine bacteria.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** Lat:33.419410120974 Lon:-111.92828937936

**Temporal Extent:** 2021 - 2022

## Methods & Sampling

In this study, one diatom and three bacteria species were grown and measured. *Minutocellus polymorphus* were incubated with or without the addition of bacteria in flasks and sampled throughout their growth (representative of bloom conditions), to determine the production of transparent exopolymeric particles (TEP) and the formation of micro-aggregates, and in roller tanks to investigate the formation of sinking aggregates (representative of end of bloom conditions). Additional details are in Cruz & Neuer, 2022.

Two growth experiments were carried out independently:

1. Growth experiment 1 with the addition of *Marinobacter adhaerens*
2. Growth experiment 2 with the addition of *Pseudoalteromonas carrageenovora* and *Vibrio thalassae*

Stock cultures of marine *Minutocellus polymorphus* (CCMP497, National Center for Marine Algae and Microbiota, NCMA) were maintained in L1 medium prepared in artificial seawater and incubated in an

environmental growth chamber (Conviron) at  $23 \pm 1$  °C. Stock cultures of *Vibrio thalassae* (DSM102810, DSMZ-German Collection of Microorganisms and Cell Cultures GmbH), *Pseudoalteromonas carrageenovora* (DSM6820, DSMZ), and *Marinobacter adhaerens* HP15 were maintained on Marine Agar (BD Difco 2216, Becton Dickinson, NJ; ZoBell, 1941) plates at  $23 \pm 1$  °C.

Triplicate cultures of *Minutocellus polymorphus* (CCMP497) with and without the addition of known particle-associated marine bacteria (*M. adhaerens*, *P. carrageenovora*, and *V. thalassae*) were sampled every other day for 19-23 days for the quantification of:

- single cells [Cell Abundance dataset]
- suspended microaggregates (aggregates ca. 5-60  $\mu\text{m}$ ) [this dataset]
- TEP (Transparent Exopolymeric Particles) [TEP Concentration dataset]

This dataset contains data on suspended microaggregates, while the other measurements can be found in the Related Datasets below.

Volume concentrations of micro-aggregates “suspended” in cultures (i.e., non-sinking particles with an equivalent spherical diameter [ESD] of 5–60  $\mu\text{m}$ ) were determined at every sampling period using a Multisizer 3 Particle Counter (Beckman Coulter, CA). Prior to fixation with glutaraldehyde, duplicate samples were diluted to a 1–10% final particle concentration with Isoton II diluent (Beckman Coulter, CA) and aggregates were measured and quantified with a 100  $\mu\text{m}$  aperture tube. The volume concentration of aggregates was calculated in  $\mu\text{m}^3$  per mL.

### Aggregation in roller tanks

To investigate the formation of visible aggregates, cultures of the nanodiatom *Minutocellus polymorphus* (CCMP497) with and without the addition of known particle-associated marine bacteria (*Marinobacter adhaerens* HP15, *Pseudoalteromonas carrageenovora*, and *Vibrio thalassae*) were incubated in cylindrical 1.25 L Plexiglass roller tanks with artificial seawater (ASW) for 7 days in the dark at 3.5 revolutions per minute (RPM) after the methodology of Shanks and Edmondson (1989).

Two sets of experiments were performed:

1. cultures were inoculated to cell concentrations simulating natural conditions in the Sargasso Sea ( $10^3$  cells per mL), and the experiment was performed with the addition of *M. adhaerens* alongside bacteria-only and diatom-only controls, as well as with the addition of microbeads
2. cultures were inoculated at higher cell concentrations ( $10^5$  cells per mL).

All treatments were performed in triplicate ( $n = 3$  tanks) in the  $10^5$  cells  $\text{mL}^{-1}$  experiments and in duplicate ( $n = 2$  tanks) in the  $10^3$  cells  $\text{mL}^{-1}$  experiments. Aggregates formed were quantified, sized, and their sinking velocities were determined. Suspended microaggregates (i.e., non-sinking particles with an equivalent spherical diameter of 5-60 microns) were quantified using a Multisizer 3 Particle Counter. TEP concentrations were determined as in Bittar et al. (2018). The stocks of Alcian-Blue dye used for TEP quantification had calibration factors (or f-factors) of 81.70 for experiments with *M. adhaerens* and 83.83 for experiments with *P. carrageenovora* and *V. thalassae*.

Only the axenic *Minutocellus*  $10^5$  experiment had hardy-enough aggregates for analysis. Please refer to the Supplemental document.

For additional Methods details, see Cruz & Neuer, 2022.

For size and velocity of sinking aggregates from roller tank experiments, see Supplemental Files section

### Data Processing Description

BCO-DMO Processing Notes:

- Excel sheet "M.polymorphus\_Microaggregates\_and\_TEP- Suspended Microaggregates.xlsx" was processed and saved as a CSV file
- added a conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- blank values in this dataset are displayed as "nd" for "no data." nd is the default missing data identifier in the BCO-DMO system.

[ [table of contents](#) | [back to top](#) ]

---

## Supplemental Files

File
<b>Cruz &amp; Neuer 2022 Roller tank aggregate sizes and velocities</b> filename: Cruz_Neuer_roller_tank_size_velocities.pdf (Portable Document Format (.pdf), 590.01 KB) MD5:edd0c5590b44ca83d5d4efd1f5226d57 Sizes and velocities of aggregates from roller tank experiments as described in Cruz & Neuer, 2022.

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Cruz, B. N., & Neuer, S. (2022). Particle-associated bacteria differentially influence the aggregation of the marine diatom *Minutocellus polymorphus*. ISME Communications, 2(1). <https://doi.org/10.1038/s43705-022-00146-z>

*Results*

Shanks, A. L., & Edmondson, E. W. (1989). Laboratory-made artificial marine snow: a biological model of the real thing. Marine Biology, 101(4), 463–470. doi:10.1007/bf00541648 <https://doi.org/10.1007/BF00541648>

*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### IsRelatedTo

Neuer, S., Cruz, B. N., Cadillo-Quiroz, H. (2022) **Laboratory data on cell abundance of *Minutocellus polymorphus* in experiments measuring TEP and production of microaggregates.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-11-04 <http://lod.bco-dmo.org/id/dataset/882570> [[view at BCO-DMO](#)]

Neuer, S., Cruz, B. N., Cadillo-Quiroz, H. (2022) **TEP concentrations of co-cultures and axenic cultures of *Minutocellus polymorphus* and particle-associated marine bacteria.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-10-25 <http://lod.bco-dmo.org/id/dataset/882581> [[view at BCO-DMO](#)]

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Experiment_ID	Experiment ID describing the experiment in which the organism grew	unitless
Culture	Experimental culture, listing the organisms grown	unitless
Culture_type	Culture type, either co-culture or axenic (diatom-only control)	unitless
Days	Days since start of experiment	days
Susp_Microaggregate_Vol_Conc_rep1	Replicate 1 volume concentration of aggregates with an equivalent spherical diameter of 5-60 microns	$10^6$ cubic microns per milliliter ( $\times 10^6 \text{ um}^3$ per mL)
Susp_Microaggregate_Vol_Conc_rep2	Replicate 2 volume concentration (times $10^6$ ) of aggregates with an equivalent spherical diameter of 5-60 microns	cubic microns per milliliter ( $\times 10^6 \text{ um}^3$ per mL)
Susp_Microaggregate_Vol_Conc_rep3	Replicate 3 volume concentration (times $10^6$ ) of aggregates with an equivalent spherical diameter of 5-60 microns	cubic microns per milliliter ( $\times 10^6 \text{ um}^3$ per mL)

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Multisizer 3 Particle Counter
<b>Generic Instrument Name</b>	Coulter Counter
<b>Dataset-specific Description</b>	Suspended microaggregates were quantified using a Multisizer 3 Particle Counter
<b>Generic Instrument Description</b>	An apparatus for counting and sizing particles suspended in electrolytes. It is used for cells, bacteria, prokaryotic cells and virus particles. A typical Coulter counter has one or more microchannels that separate two chambers containing electrolyte solutions. from <a href="https://en.wikipedia.org/wiki/Coulter_counter">https://en.wikipedia.org/wiki/Coulter_counter</a>

<b>Dataset-specific Instrument Name</b>	Epifluorescence microscope
<b>Generic Instrument Name</b>	Fluorescence Microscope
<b>Dataset-specific Description</b>	Single cell abundance of diatoms and bacteria were determined using epifluorescence microscopy
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of fluorescence and phosphorescence instead of, or in addition to, reflection and absorption of visible light. Includes conventional and inverted instruments.

<b>Dataset-specific Instrument Name</b>	Roller tank
<b>Generic Instrument Name</b>	Roller Tank
<b>Dataset-specific Description</b>	Cultures were incubated in 1.25 L roller tanks for 7 days in the dark at 3.5 revolutions per minute (RPM)
<b>Generic Instrument Description</b>	Rolling tanks, which keep particles in suspension, thus simulating aggregate formation in situ. Marine snow experiments are conducted in roller tanks, which turn continuously, keeping marine snow in suspension. It is important for marine snow not to touch surfaces. The rolling tanks, which keep particles in suspension, thus simulate aggregate formation in situ. Marine snow formation due to different types of oil was tested. Some treatments are easily identifiable as containing oil by their color (middle). UCSB, CA 2012.

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### Aggregation of Marine Picoplankton (Marine Plankton Aggregation)

**Coverage:** Bermuda Atlantic Time-Series station

NSF abstract:

Marine phytoplankton are microscopic algae that live in the sunlit zone of the ocean. They play an important role in the uptake of carbon dioxide from the atmosphere through photosynthesis, similar to what plants do on land, and are the basis of the marine food web. However, instead of storing this organic carbon in leaf tissue and roots, marine phytoplankton are grazed by planktonic animals, or die and subsequently sink out of the sunlit zone in the form of aggregates, also called "Marine Snow". These particles not only export the organic carbon contained in their cells to the deep ocean, but also serve as food for animals and bacteria that live in the deep. A considerable portion of these phytoplankton are extremely small, among the tiniest of all organisms known. These extremely small cells have not been thought to play an important role in the formation and sinking of marine snow; however, recent findings challenge this view. This project will investigate how the smallest of these phytoplankton contribute to the rain of sinking particles from the sunlit surface to the deep ocean. This research is important because, in some of the largest expanses of the open oceans, these minute cells dominate the phytoplankton community, and larger plankton organisms are very sparse. The project, through a combination of work in the laboratory and at a field station, will shed light on how these tiny phytoplankton cells make aggregates, which ultimately enable them to sink as "Marine Snow". The project also provides unique opportunities for undergraduate students at Arizona State University, a land-locked public university, to gain experience in working with marine research. The project will serve to educate one PhD student, one MS student in an accelerated BS-MS program, and 8-10 undergraduate students/semester in a unique, inquiry based learning effort termed Microbial Education Training and Outreach (MENTOR). The undergraduate students will also participate in Arizona State University (ASU)'s School of Life Sciences, Undergraduate Research Program (SOLUR), which seeks to increase the participation of minorities in science. They will also contribute towards developing web and classroom materials, based on this project, which will then be distributed through a partnership with the award-winning ASU-sponsored Ask A Biologist K-12 web site.

The oceanic "biological carbon pump", the photosynthetically mediated transformation of dissolved inorganic carbon into particulate and dissolved organic carbon and its subsequent export to deep water, functions as a significant driver of atmospheric carbon uptake by the oceans. The traditional view of the biological carbon pump in the ocean is that of sinking of large aggregates (marine snow) or fecal pellets, which are made up of large, mineral ballasted cells of phytoplankton. However, recent evidence, stemming from in situ investigations of particulate matter, trap studies and modelling studies, have shown that micron-sized phytoplankton such as picocyanobacteria as well as picoeukaryotes can contribute significantly to the sinking of particulate matter.

The specific mechanisms behind the sinking of these micrometer sized cells remain elusive as the cells are too small to sink on their own, and mesozooplankton is likely unable to ingest single cells. Intriguingly, recent research by the investigators has shown that the ubiquitous picocyanobacteria *Synechococcus* are able to form aggregates and sink at velocities comparable to those of marine snow. They found that the matrix of the *Synechococcus* aggregates was made of Transparent Exopolymeric Particles (TEP), and that TEP production was enhanced under nutrient limited culture conditions. Interaction with clays and presence of heterotrophic bacteria also enhanced aggregation and sinking velocity. This study aims to further investigate aggregation of other common picoplankton in the laboratory and aggregation occurring in natural settings at an oligotrophic open ocean site, the Bermuda Atlantic Time-series Site (BATS). Ultimately, this project will increase and refine our understanding of the role of the smallest phytoplankton in aggregation and sinking - information vital to understanding carbon cycling processes in the oceans.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1658527</a>

[ [table of contents](#) | [back to top](#) ]