

# Kelp understory blade growth data from macroalgal surveys in Sitka Sound, Alaska kelp beds from 2018 to 2019

**Website:** <https://www.bco-dmo.org/dataset/882043>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2022-10-10

## Project

» [CAREER: Energy fluxes and community stability in a dynamic, high-latitude kelp ecosystem](#) (High latitude kelp dynamics)

Contributors	Affiliation	Role
<a href="#">Kroeker, Kristy J.</a>	University of California-Santa Cruz (UCSC)	Principal Investigator
<a href="#">Bell, Lauren E.</a>	University of California-Santa Cruz (UCSC)	Student, Contact
<a href="#">York, Amber D.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Kelp understory blade growth data from macroalgal surveys in Sitka Sound, Alaska kelp beds from 2018 to 2019.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Methods & Sampling](#)
    - [Data Processing Description](#)
  - [Data Files](#)
  - [Supplemental Files](#)
  - [Related Publications](#)
  - [Parameters](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** N:57.039 E:-135.278 S:56.9875 W:-135.357

**Temporal Extent:** 2018-07-20 - 2019-07-22

## Methods & Sampling

We monitored monthly growth and loss of dominant kelp species in Sitka Sound from January 2017 - February 2018 at Breast Is. and Harris Is. (*M. pyrifera* only), and from July 2018 - July 2019 at Harris Is., Breast Is., and Samsing Pinnacle (*M. pyrifera*, *N. fimbriatum*, *H. nigripes*). At each site, we identified 12 - 15 "adult" individuals of each species (*M. pyrifera*: at least one frond reaching the surface; *N. fimbriatum* and *H. nigripes*: maximum blade length > 20 cm) along a 5 - 6 m depth (MLLW) contour with numbered tags. Each month, we re-surveyed tagged *M. pyrifera* plants for frond density, with zip ties loosely bound around new fronds exceeding 1 m in height to distinguish new growth. For tagged *N. fimbriatum* and *H. nigripes* plants, each month we punched a new hole through the thallus at 10 cm from the intercalary meristem (Parke 1948), and we measured blade morphometrics (maximum blade length and width) and distance from meristem to the previous month's punched hole. When previously tagged individuals were not re-sighted after two consecutive months, we assumed they had been physically removed from the substrate, either through grazing or abiotic factors.

Additional Funding Details:

In addition to primary funding from the NSF award OCE-1752600 additional funding was provided from The

David and Lucile Packard Foundation and the North Pacific Research Board's Graduate Student Research Award (1748-01) to Lauren Bell, PhD University of California Santa Cruz, Award title: "Fish Habitat, Fishes and Invertebrates, Lower Trophic Level Productivity Effect of substrate on herring roe response to global change."

## Data Processing Description

BCO-DMO Data Manager Processing Notes:

\* File "HNIG.NFIM\_blade.growth.csv" imported into the BCO-DMO data system.

\* Date format changed to ISO format

\* Species list with codes and scientific names extracted from parameter information. Matched to known taxon ids using WoRMS taxa match (2022-09-06). The spelling of "Laminaria setchellii" changed to "Laminaria setchellii" with two Ls after confirming the change with the data submitter. Species list along with identifiers attached as a supplemental data table.

\* Latitude and Longitude added to main data table from the provided site list.

[ [table of contents](#) | [back to top](#) ]

---

## Data Files

File
<b>blade_growth.csv</b> (Comma Separated Values (.csv), 36.55 KB) MD5:c509e473841189a2ff63d1fbbde873f6
Primary data file for dataset ID 882043

[ [table of contents](#) | [back to top](#) ]

---

## Supplemental Files

File
<b>Sitka macroalgal survey site list</b> filename: site_list.csv (Comma Separated Values (.csv), 259 bytes) MD5:785b37ebc9f99bea71257234529ad278
Site list for macroalgal surveys conducted in Sitka, Alaska between 2017 to 2020.
Parameters (column name, description, units): Site, Site name,unitless Latitude, latitude of site, decimal degrees Longitude,longitude of site, decimal degrees
<b>Sitka macroalgal survey species list</b> filename: species_list.csv (Comma Separated Values (.csv), 490 bytes) MD5:983f490f673acb204083528f4f11e380
Species list for macroalgal surveys conducted in Sitka, Alaska between 2017 to 2020.
Parameters (column name, description, units): Sp, species code used in related datasets (e.g. MPYR),unitless ScientificName,The accepted scientific name for the species (as of 2022-09),unitless AphiaID,Taxonomic identifier AphiaID for the species (see World Register of Marine Species),unitless LSID,Life Sciences Identifier (LSID) for the species,unitless

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Bell, L. E., & Kroeker, K. J. (2022). Standing Crop, Turnover, and Production Dynamics of *Macrocystis pyrifera* and Understory Species *Hedophyllum nigripes* and *Neoagarum fimbriatum* in High Latitude Giant Kelp Forests. *Journal of Phycology*, 58(6), 773–788. Portico. <https://doi.org/10.1111/jpy.13291>  
*Results*

Parke, M. (1948). Studies on British Laminariaceae. I. Growth in *Laminaria Saccharina* (L.) Lamour. *Journal of the Marine Biological Association of the United Kingdom*, 27(3), 651–709.  
<https://doi.org/10.1017/s0025315400056071> <https://doi.org/10.1017/S0025315400056071>  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Site	Name of rocky reef location in Sitka Sound where survey occurred.	unitless
Latitude	Site latitude	decimal degrees
Longitude	Site longitude	decimal degrees
Transect	Identity of recurring 30 x 2 m transect in which surveyed plants were located (C-D)	unitless
plant_ID	Alpha-numeric identifier for each tagged alga. Unique by site.	unitless
date_T0	Date of survey period start (T0) in ISO 8601 format (YYYY-MM-DD)	unitless
date_T1	Date of survey period end (T1) in ISO 8601 format (YYYY-MM-DD)	unitless
dataquality_flag	Data quality identifier. [G = Good = Punched hole resighted without issue. F = fair = some grazing or deterioration of blade, but still trust growth data. P = poor = difficult to make out hole, data in question. Where hole enlarged by grazing and diver noted a possible range of where it could be, used conservative estimate (minimum growth), and noted as F]	unitless
Sp	Species of understory macroalgae surveyed (NFIM = <i>Neoagarum fimbriatum</i> , HNIG = <i>Hedophyllum nigripes</i> )	unitless
maxlength_T0	Maximum blade length at survey period start (T0). Detection limit: 1cm	centimeters (cm)
maxwidth_T0	Maximum blade width at survey period start (T0). Detection limit: 1cm	centimeters (cm)
stipetohole_T0	Distance from intercalary meristem to punched hole at survey period start (T0). Detection limit: 0.1cm	centimeters (cm)
maxlength_T1	Maximum blade length at survey period end (T1). Detection limit: 1cm	centimeters (cm)
maxwidth_T1	Maximum blade width at survey period end (T1). Detection limit: 1cm	centimeters (cm)
stipetohole_T1	Distance from intercalary meristem to punched hole at survey period end (T1). Detection limit: 0.1cm	centimeters (cm)
Interval_days	Number of days elapsed between T0 and T1. Detection limit: 1 day	days
holepastend_flag	Data quality identifier. Was punched hole missing at end of survey period (due to grazing or erosion of blade)? Y = yes, N = no	unitless

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

**CAREER: Energy fluxes and community stability in a dynamic, high-latitude kelp ecosystem (High latitude kelp dynamics)**

**Coverage:** SE Alaskan coastal waters

### *NSF Award Abstract:*

High latitude kelp forests support a wealth of ecologically and economically important species, buffer coastlines from high-energy storms, and play a critical role in the marine carbon cycle by sequestering and storing large amounts of carbon. Understanding how energy fluxes and consumer-resource interactions vary in these kelp communities is critical for defining robust management strategies that help maintain these valuable ecosystem services. In this integrated research and education program, the project team will investigate how consumer populations respond to variability in temperature, carbonate chemistry and resource quality to influence the food webs and ecosystem stability of kelp forests. A comprehensive suite of studies conducted at the northern range limit for giant kelp (*Macrocystis pyrifera*) in SE Alaska will examine how kelp communities respond to variable environmental conditions arising from seasonal variability and changing ocean temperature and acidification conditions. As part of this project, undergraduate and high school students will receive comprehensive training through (1) an immersive field-based class in Sitka Sound, Alaska, (2) intensive, mentored research internships, and (3) experiential training in science communication and public outreach that will include a variety of opportunities to disseminate research findings through podcasts, public lectures and radio broadcasts.

Consumer-resource interactions structure food webs and govern ecosystem stability, yet our understanding of how these important interactions may change under future climatic conditions is hampered by the complexity of direct and indirect effects of multiple stressors within and between trophic levels. For example, environmentally mediated changes in nutritional quality and chemical deterrence of primary producers have the potential to alter herbivory rates and energy fluxes between primary producers and consumers, with implications for ecosystem stability. Moreover, the effects of global change on primary producers are likely to depend on other limiting resources, such as light and nutrients, which vary seasonally in dynamic, temperate and high latitude ecosystems. In marine ecosystems at high latitude, climate models predict that ocean acidification will be most pronounced during the winter months, when primary production is limited by light. This project is built around the hypothesis that there could be a mismatch in the energetic demands of primary consumers caused by warming and ocean acidification and resource availability and quality during winter months, with cascading effects on trophic structure and ecosystem stability in the future. Through complementary lab and field experiments, the project team will determine 1) how temperature and carbonate chemistry combine to affect primary consumer bioenergetics across a diversity of species and 2) the indirect effects of ocean acidification and warming on primary consumers via environmentally mediated changes in the availability, nutritional quality and palatability of primary producers across seasons. Using the data from the laboratory and field experiments, the project team will 3) construct a model of the emergent effects of warming and ocean acidification on trophic structure and ecosystem stability in seasonally dynamic, high latitude environments.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1752600</a>

[ [table of contents](#) | [back to top](#) ]