

Field physiochemical parameters including nutrient concentrations and nitrogen specific uptake rates from samples collected between 2017 and 2019 from the Arctic Ocean, California Coastal Current, and a Chesapeake Bay estuary

Website: <https://www.bco-dmo.org/dataset/896181>

Data Type: Cruise Results

Version: 1

Version Date: 2023-05-17

Project

» [Collaborative Research: Creatine Cycling in Marine Bacterial and Phytoplankton Assemblages](#) (Creatine Cycling)

Contributors	Affiliation	Role
Bronk, Deborah A.	Virginia Institute of Marine Science (VIMS)	Principal Investigator, Contact
Wawrik, Boris	University of Oklahoma (OU)	Co-Principal Investigator
Yang, Zhibo	University of Oklahoma (OU)	Co-Principal Investigator
Stanley, Brianna	Virginia Institute of Marine Science (VIMS)	Scientist
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Nitrogen is an important macronutrient for biological growth and while research has been traditionally focused on the dissolved inorganic nitrogen pool, dissolved organic nitrogen (DON) is an important, but historically less understood source of nitrogen. This work focused on creatine, a DON metabolite recognized in human nutrition that has not been heavily studied in aquatic systems. Our research investigates the extent and rate at which aquatic organisms may uptake (use) creatine and compares those rates to other nitrogen compounds including ammonium, nitrate, urea, and amino acids. Samples were collected between 2017 and 2019 and cover three North American regions. These include the polar Arctic Ocean, the California Coastal Current (Baja), and a Chesapeake Bay estuary (York River). Included in this dataset are ambient physiochemical parameters for these systems, along with measured specific uptake rates.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:72.55 E:-76.687 S:28.29 W:-115.91

Temporal Extent: 2017-05-03 - 2019-07-16

Methods & Sampling

Site water for incubation experiments and for the analysis of ambient nutrients was collected slightly differently for each of the sampling regions. In the Arctic, site water for incubation experiments was collected on the R/V Ukpik and R/V Sikuliaq (cruise ID number SK201712S) during the summer of 2017 using a submersible pump and CTD rosette, respectively. In Baja, sampling occurred twice on the R/V Robert Gordon Sproul during both May and October 2017 (cruise ID numbers SP1714 and SP1727), and site water was collected using a CTD rosette (Turk-Kubo et al. 2021). In Virginia, surface sites in the York River were sampled by directly filling an acid-washed PETG carboy over the side. Filtrations for sample filtrate and filters also varied. These York Rivers samples were obtained during day trips aboard a variety of small powerboats (<21 feet), every other month from June 2018 to July 2019. Powerboats are housed out of the Virginia Institute of Marine Science.

In the Arctic, filtrations were conducted in parallel, which separately passed site water through a larger Nucleopore Membrane filter (3.0 micrometer (μm) nominal pore size) and a smaller Whatman GF-75 filter (nominal pore size 0.3 μm). This same combination of filters was used in Baja in October 2017. In May 2017, Baja sampling used the same filters, but filtered sequentially, meaning that site water was first passed through the Nucleopore filter before passing through the GF-75. The York River site samples were only filtered using the GF-75. Filtrate for all regions was collected from the GF-75 filtration and kept for later nutrient analyses and filters

were used for chlorophyll a analysis.

Pigments (chlorophyll a and phaeopigments) were measured after extraction with 90% acetone overnight (Parsons et al. 1984; Arar and Collins 1997). Concentrations of ammonium were analyzed using the Koroleff (1983) method and amino acids were measured as dissolved primary amines (DPA; Parsons et al. 1984). Concentrations of nitrate, nitrite, phosphate, and silica were measured using a Lachat 8500 Quickchem autoanalyzer. Urea was analyzed using the monoxime method (Price and Harrison 1987). Concentrations of total dissolved nitrogen and dissolved organic carbon were assessed using a Shimadzu TOC-V TNM (Hansell 1993). Dissolved organic nitrogen is calculated as the difference between total dissolved nitrogen and inorganic nitrogen.

Nitrogen uptakes were measured after incubation for a set time (1 to 24 hours) in either 0.5 or 1-liter PETG bottles. Stable isotope tracer methods were used according to those described in Baer et al (2017). Uptake rates for inorganic and organic nitrogen were measured by incubating water with ^{15}N - ammonium, nitrate, creatine, urea, and/or an amino acid mixture under *in situ* light and temperature conditions in a flow-through system on deck or in a cold room set to ambient site water temperatures. Rate incubation experiments were terminated with the same filtration methods as collection of site water ambients with the exception that a Sterlitech silver filter was used in lieu of the Nucleopore Membrane filter. The exact combination of nitrogen substrates varied between sites and regions. All nitrogen uptake samples were analyzed on a Sercon Integra 2 isotope ratio mass spectrometer.

Known Issues/Problems:

Note that for the October sampling in the Baja region, chlorophyll a concentrations for the nutrient size fractions are unavailable. Instead, chlorophyll a concentrations for the $>0.7\ \mu\text{m}$ size fraction are available thanks to the Arrigo and Zehr dataset (listed under "Related Datasets").

Data Processing Description

Data Processing:

It is important to note that potential specific uptake rates are reported based on sampled size fraction (i.e. >3.0 or $>0.3\ \mu\text{m}$). When applicable, the rates of different size fractions were combined for the final reported specific uptake rate (i.e. >3.0 and $0.3 - 3.0\ \mu\text{m}$ fractions added to report a $>0.3\ \mu\text{m}$ rate). Rates of ammonium-specific uptake were corrected for isotope dilution when possible. These rates were corrected according to Gilbert et al. (1982).

BCO-DMO Processing:

- removed "NA" and "ND" (missing data values are blank/empty in the final .csv data file);
- converted the Date column to YYYY-mm-dd format;
- converted the Time column to HH:MM format;
- created the ISO_DateTime_UTC column, which presents date and time in UTC in ISO 8601 format;
- renamed fields to comply with BCO-DMO naming conventions in which the only allowed characters are letters (A-Z, a-z), numbers (0-9), and underscores.

[[table of contents](#) | [back to top](#)]

Data Files

File
creatine_nutrients_n_rates.csv (Octet Stream, 26.24 KB) MD5:8ceb2a679904acfff535b19228a77d46
Primary data file for dataset ID 896181

[[table of contents](#) | [back to top](#)]

Related Publications

Arar, E. J. AND G. B. COLLINS. Method 445.0 In Vitro Determination of Chlorophyll a and Pheophytin in Marine and Freshwater Algae by Fluorescence. U.S. Environmental Protection Agency, Washington, DC, 1997.

https://cfpub.epa.gov/si/si_public_record_report.cfm?Lab=NERL&dirEntryId=309417

Methods

Baer, S. E., Sipler, R. E., Roberts, Q. N., Yager, P. L., Frischer, M. E., & Bronk, D. A. (2017). Seasonal nitrogen uptake and regeneration in the western coastal Arctic. *Limnology and Oceanography*, 62(6), 2463-2479. doi:[10.1002/lno.10580](https://doi.org/10.1002/lno.10580)

Methods

Gilbert, P. M., Lipschultz, F., McCarthy, J. J., & Altabet, M. A. (1982). Isotope dilution models of uptake and remineralization of ammonium by marine plankton. *Limnology and Oceanography*, 27(4), 639-650. Portico. <https://doi.org/10.4319/lno.1982.27.4.0639>

Methods

Hansell, D. A. (1993). Results and observations from the measurement of DOC and DON in seawater using a high-temperature catalytic oxidation technique. *Marine Chemistry*, 41(1-3), 195-202. [https://doi.org/10.1016/0304-4203\(93\)90119-9](https://doi.org/10.1016/0304-4203(93)90119-9)

Methods

Koroleff, F. (1983). Simultaneous Oxidation of Nitrogen and Phosphorus Compounds by Persulfate. In K. Grasshoff, M. Eberhardt, and F. Kremling [eds.], *Methods of Seawater Analysis*. GMB: Verlag Chemie.

Methods

Parsons, T. R., Maita, Y., & Lalli, C.M. (1984). A manual of chemical and biological methods for seawater analysis. Pergamon Press. doi:10.1016/c2009-0-07774-5 <https://doi.org/10.1016/C2009-0-07774-5>
Methods

Price, N. M., & Harrison, P. J. (1987). Comparison of methods for the analysis of dissolved urea in seawater. *Marine Biology*, 94(2), 307-317. doi:10.1007/bf00392945 <https://doi.org/10.1007/BF00392945>
Methods

Turk-Kubo, K. A., Mills, M. M., Arrigo, K. R., van Dijken, G., Henke, B. A., Stewart, B., Wilson, S. T., & Zehr, J. P. (2021). UCYN-A/haptophyte symbioses dominate N₂ fixation in the Southern California Current System. *ISME Communications*, 1(1). <https://doi.org/10.1038/s43705-021-00039-7>
Results

[[table of contents](#) | [back to top](#)]

Related Datasets

IsRelatedTo

Arrigo, K. R., Zehr, J. P. (2023) **CTD sensor data from two cruises from R/V Robert Gordon Sproul SP1714 in the California Current waters off the coast of Southern California and Baja California from 2017-2018**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2023-11-29 doi:10.26008/1912/bco-dmo.774459.1 [[view at BCO-DMO](#)]

Relationship Description: Additional data from the SP1714 and SP1727 cruises in the Baja region.

Sipler, R. & Bronk, D. (2021). Nutrients, nitrogen fixation, nitrogen uptake and carbon uptake data collected from the Western Arctic Ocean, 2016-2017. Arctic Data Center. urn:uuid:2017c57b-07a7-4e25-98a6-d2954aa59bb6.

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Ship	Name of research vessel	unitless
Region	General geographic region for the data collected	unitless
Designator	Numerical cruise number or other descriptor	unitless
Site_Name	Name or descriptor of sampled site location	unitless
Latitude	Latitude (the North direction is represented by positive numbers)	decimal degrees
Longitude	Longitude (the West direction is represented by negative numbers)	decimal degrees
Date	Date of sample collection in the local time zone	unitless
Time	Time of sample collection recorded in the respective local time using a 24 hour clock	unitless
Time_Zone	Respective local time zone where AKDT is Alaska Daylight Time, EST is Eastern Standard Time, and EDT is Eastern Daylight Time	unitless
ISO_DateTime_UTC	Date and time of sample collection in UTC in ISO 8601 format	unitless
Water_Depth	General descriptor (e.g., surface, Chlorophyll maximum, etc.)	unitless
Sample_Depth	Depth sample collected at in meters	meters (m)
Total_Water_Column_Depth	Depth of the total water column in meters	meters (m)
Salinity	Salinity of site water in parts per trillion	parts per trillion
Temperature	Temperature of site water in degrees Celsius	degrees Celsius
Total_Dissolved_N	Total dissolved nitrogen in micromoles nitrogen per liter	micromoles per liter (umol/L)
Total_Dissolved_Nz_Std	One standard deviation of Total_Dissolved_N	micromoles per liter (umol/L)

Nitrite	Nitrite in micromoles nitrogen per liter; 0.000 = Below detection; Detection limit: 0.05 umol N L-1	micromoles per liter (umol/L)
Nitrite_Std	One standard deviation of Nitrite. "SL" means "Sample lost".	micromoles per liter (umol/L)
Nitrate	Nitrate in micromoles nitrogen per liter; 0.000 = Below detection; Detection limit: 0.05 umol N L-1	micromoles per liter (umol/L)
Nitrate_Std	One standard deviation of Nitrate	micromoles per liter (umol/L)
Ammonium	Ammonium in micromoles nitrogen per liter; 0.000 = Below detection; Detection limit: 0.03 umol N L-1	micromoles per liter (umol/L)
Ammonium_Std	One standard deviation of Ammonium	micromoles per liter (umol/L)
Dissolved_Organic_N	Dissolved organic nitrogen in micromoles nitrogen per liter	micromoles per liter (umol/L)
Dissolved_Organic_N_Std	One standard deviation of Dissolved_Organic_N. "SL" means "Sample lost".	micromoles per liter (umol/L)
Urea	Urea in micromoles nitrogen per liter; 0.000 = Below detection; Detection limit: 0.1 umol N L-1	micromoles per liter (umol/L)
Urea_Std	One standard deviation of Urea	micromoles per liter (umol/L)
Dissolved_Primary_Amines	Dissolved primary amine in micromoles nitrogen per liter; 0.000 = Below detection; Detection limit: 0.025 umol N L-1	micromoles per liter (umol/L)
Dissolved_Primary_Amines_Std	One standard deviation of Dissolved_Primary_Amines	micromoles per liter (umol/L)
Dissolved_Organic_C	Dissolved organic carbon in micromoles carbon per liter	micromoles per liter (umol/L)
Dissolved_Organic_C_Std	One standard deviation of Dissolved_Organic_C	micromoles per liter (umol/L)
Phosphate	Phosphate in micromoles phosphorus per liter; 0.000 = Below detection; Detection limit: 0.05 umol N L-1	micromoles per liter (umol/L)
Phosphate_Std	One standard deviation of Phosphate	micromoles per liter (umol/L)
Silicate	Silicate in micromoles silica per liter; 0.000 = Below detection; Detection limit: 0.11 umol N L-1	micromoles per liter (umol/L)
Silicate_Std	One standard deviation of Silicate	micromoles per liter (umol/L)
Chl_a_gt_3um	Chlorophyl a in micrograms per liter in the size fraction representing the community greater than 3.0 micrometers (>3.0um); 0.000 = Below detection; Detection limit: 0.025 ug L-1	micrograms per liter (ug/L)
Chl_a_gt_3um_Std	One standard deviation of Chl_a_gt_3um	micrograms per liter (ug/L)

Chl_a_gt_7tenths_um	Chlorophyl a in micrograms per liter in the size fraction representing the community greater than 0.7 micrometers (>0.7um); 0.000 = Below detection; Detection limit: 0.025 ug L-1	micrograms per liter (ug/L)
Chl_a_gt_7tenths_um_Std	One standard deviation of Chl_a_gt_7tenths_um	micrograms per liter (ug/L)
Phaeo_gt_3um	Phaeopigment in micrograms per liter in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	micrograms per liter (ug/L)
Phaeo_gt_3um_Std	One standard deviation of Phaeo_gt_3um	micrograms per liter (ug/L)
Chl_a_gt_3tenths_um	Chlorophyl a in micrograms per liter in the size fraction representing the community greater than 0.3 micrometers (>0.3um); 0.000 = Below detection; Detection limit: 0.025 ug L-1	micrograms per liter (ug/L)
Chl_a_gt_3tenths_um_Std	One standard deviation of Chl_a_gt_3tenths_um	micrograms per liter (ug/L)
Phaeo_gt_3tenths_um	Phaeopigment in micrograms per liter in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	micrograms per liter (ug/L)
Phaeo_gt_3tenths_um_Std	One standard deviation of Phaeo_gt_3tenths_um	micrograms per liter (ug/L)
Creatine_V_gt_3um	Specific uptake rate of creatine per hour in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
Creatine_V_gt_3um_Std	One standard deviation of Creatine_V_gt_3um	per hour
Creatine_V_3tenths_to_3um	Specific uptake rate of creatine per hour in the size fraction representing the community between 0.3 and 3.0 micrometers (0.3-3.0um)	per hour
Creatine_V_3tenths_to_3um_Std	One standard deviation of Creatine_V_3tenths_to_3um	per hour
Creatine_V_gt_3tenths_um	Specific uptake rate of creatine per hour in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
Creatine_V_gt_3tenths_um_Std	One standard deviation of Creatine_V_gt_3tenths_um	per hour
Isotope_Dilution_Corrected_Ammonium_V_gt_3um	Specific uptake rate per hour for ammonium, corrected for isotope dilution, in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
Isotope_Dilution_Corrected_Ammonium_V_gt_3um_Std	One standard deviation of Isotope_Dilution_Corrected_Ammonium_V_gt_3um	per hour
Isotope_Dilution_Corrected_Ammonium_V_gt_3tenths_um	Specific uptake rate per hour for ammonium, corrected for isotope dilution, in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
Isotope_Dilution_Corrected_Ammonium_V_gt_3tenths_um_Std	One standard deviation of Isotope_Dilution_Corrected_Ammonium_V_gt_3tenths_um	per hour
Ammonium_V_gt_3um	Specific uptake rate of ammonium per hour in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
Ammonium_V_gt_3um_Std	One standard deviation of Ammonium_V_gt_3um	per hour
Ammonium_V_3tenths_to_3um	Specific uptake rate of ammonium per hour in the size fraction representing the community between 0.3 and 3.0 micrometers (0.3-3.0um)	per hour
Ammonium_V_3tenths_to_3um_Std	One standard deviation of Ammonium_V_3tenths_to_3um	per hour
Ammonium_V_gt_3tenths_um	Specific uptake rate of ammonium per hour in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
Ammonium_V_gt_3tenths_um_Std	One standard deviation of Ammonium_V_gt_3tenths_um	per hour

Nitrate_V_gt_3um	Specific uptake rate of nitrate per hour in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
Nitrate_V_gt_3um_Std	One standard deviation of Nitrate_V_gt_3um	per hour
Nitrate_V_gt_3tenths_um	Specific uptake rate of nitrate per hour in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
Nitrate_V_gt_3tenths_um_Std	One standard deviation of Nitrate_V_gt_3tenths_um	per hour
Urea_V_gt_3um	Specific uptake rate of urea per hour in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
Urea_V_gt_3um_Std	One standard deviation of Urea_V_gt_3um	per hour
Urea_V_gt_3tenths_um	Specific uptake rate of urea per hour in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
Urea_V_gt_3tenths_um_Std	One standard deviation of Urea_V_gt_3tenths_um	per hour
AA_V_gt_3um	Specific uptake rate of amino acids per hour in the size fraction representing the community greater than 3.0 micrometers (>3.0um)	per hour
AA_V_gt_3um_Std	One standard deviation of AA_V_gt_3um	per hour
AA_V_3tenths_to_3um	Specific uptake rate of amino acids per hour in the size fraction representing the community between 0.3 and 3.0 micrometers (0.3-3.0um)	per hour
AA_V_3tenths_to_3um_Std	One standard deviation of AA_V_3tenths_to_3um	per hour
AA_V_gt_3tenths_um	Specific uptake rate of amino acids per hour in the size fraction representing the community greater than 0.3 micrometers (>0.3um)	per hour
AA_V_gt_3tenths_um_Std	One standard deviation of AA_V_gt_3tenths_um	per hour

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Sercon Integra2 Mass Spectrometer
Generic Instrument Name	Mass Spectrometer
Dataset-specific Description	Used to measure isotope ratios and particulate masses
Generic Instrument Description	General term for instruments used to measure the mass-to-charge ratio of ions; generally used to find the composition of a sample by generating a mass spectrum representing the masses of sample components.

Dataset-specific Instrument Name	8-liter Niskin bottles
Generic Instrument Name	Niskin bottle
Dataset-specific Description	Used for site water sampling on R/V Sikuliaq and R/V Robert Gordon Sproul cruises
Generic Instrument Description	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

Dataset-specific Instrument Name	Lachat QuickChem 8500 autoanalyzer
Generic Instrument Name	Nutrient Autoanalyzer
Dataset-specific Description	Used to measure nitrate, nitrite, phosphate, and silicate concentrations
Generic Instrument Description	Nutrient Autoanalyzer is a generic term used when specific type, make and model were not specified. In general, a Nutrient Autoanalyzer is an automated flow-thru system for doing nutrient analysis (nitrate, ammonium, orthophosphate, and silicate) on seawater samples.

Dataset-specific Instrument Name	Johnson Pump model # 16004
Generic Instrument Name	Pump
Dataset-specific Description	Used to collect site water samples on R/V Ukpik cruise
Generic Instrument Description	A pump is a device that moves fluids (liquids or gases), or sometimes slurries, by mechanical action. Pumps can be classified into three major groups according to the method they use to move the fluid: direct lift, displacement, and gravity pumps

Dataset-specific Instrument Name	Shimadzu 5000A TOC-V/TNM
Generic Instrument Name	Shimadzu TOC-V Analyzer
Dataset-specific Description	Used to measure dissolved organic carbon and total dissolved nitrogen concentrations
Generic Instrument Description	A Shimadzu TOC-V Analyzer measures DOC by high temperature combustion method.

Dataset-specific Instrument Name	Shimadzu RF-600 Spectrofluorophotometer
Generic Instrument Name	Spectrophotometer
Dataset-specific Description	Used to measure concentrations of dissolved primary amines
Generic Instrument Description	An instrument used to measure the relative absorption of electromagnetic radiation of different wavelengths in the near infra-red, visible and ultraviolet wavebands by samples.

Dataset-specific Instrument Name	Turner Design Model 10-AU fluorometer
Generic Instrument Name	Turner Designs Fluorometer 10-AU
Dataset-specific Description	Used to measure chlorophyll a and phaeopigment concentrations
Generic Instrument Description	The Turner Designs 10-AU Field Fluorometer is used to measure Chlorophyll fluorescence. The 10AU Fluorometer can be set up for continuous-flow monitoring or discrete sample analyses. A variety of compounds can be measured using application-specific optical filters available from the manufacturer. (read more from Turner Designs, turnerdesigns.com, Sunnyvale, CA, USA)

Dataset-specific Instrument Name	Shimadzu UV-1800 spectrophotometer
Generic Instrument Name	UV Spectrophotometer-Shimadzu
Dataset-specific Description	Used to measure urea and ammonium concentrations
Generic Instrument Description	The Shimadzu UV Spectrophotometer is manufactured by Shimadzu Scientific Instruments (ssi.shimadzu.com). Shimadzu manufacturers several models of spectrophotometer; refer to dataset for make/model information.

[[table of contents](#) | [back to top](#)]

Deployments

Ukpik July-August 2017

Website	https://www.bco-dmo.org/deployment/896205
Platform	R/V Ukpik
Start Date	2017-07-25
End Date	2017-08-03
Description	R/V Ukpik July 25 - August 3, 2017 Short day to overnight trips Chief Scientist - Rachel Sipler (rsipler@bigelow.org)

SKQ201712S

Website	https://www.bco-dmo.org/deployment/896208
Platform	R/V Sikuliaq
Start Date	2017-08-06
End Date	2017-08-22
Description	R/V Sikuliaq SK201712S August 6 - August 22, 2017 Chief Scientist - Lauren Juranek (laurie.juranek@oregonstate.edu) See more cruise details at R2R: https://www.rvdata.us/search/cruise/SKQ201712S

SP1714

Website	https://www.bco-dmo.org/deployment/699986
Platform	R/V Robert Gordon Sproul
Start Date	2017-05-03
End Date	2017-05-11
Description	R/V Robert Gordon Sproul Cruise SP1714 May 3 - 11, 2017 Chief Scientist - Matthew Mills (mmmills@stanford.edu) See more cruise information from R2R: https://www.rvdata.us/search/cruise/SP1714

SP1727

Website	https://www.bco-dmo.org/deployment/774496
Platform	R/V Robert Gordon Sproul
Start Date	2017-10-04
End Date	2017-10-11
Description	R/V Robert Gordon Sproul Cruises SP1727 October 4 - 11, 2017 Chief Scientist - Matthew Mills (mmmills@stanford.edu) See more cruise information from R2R: https://www.rvdata.us/search/cruise/SP1727

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Creatine Cycling in Marine Bacterial and Phytoplankton Assemblages (Creatine Cycling)

Coverage: Atlantic bight

NSF Award Abstract:

High rates of dissolved organic nitrogen (DON) production and utilization in aquatic systems are typically attributed to microbial activity. Though it is known that there is a tight coupling between the production and consumption of biologically available DON, the composition, dynamics, and ecological significance of this rapidly cycled DON pool are less well understood. This proposal focuses on a component of the DON pool, creatine, which is historically understood as a product of metazoan activity, but appears to be both produced by phytoplankton and consumed by marine bacteria. Creatine is present in seawater in measurable quantities, which led to the hypothesis that creatine may be a significant component of the marine DON cycle. DON cycling likely has a bearing on fundamental marine ecosystem processes with large implications for carbon and nitrogen turnover on a global scale. Broader impacts of this project will include outreach that focuses on connecting scientists with K-12 students through research experiences for teachers and lesson development in collaboration with the K20 Center for Educational and Community Renewal, a statewide education research and development center at the University of Oklahoma. The project will integrate the research with inquiry-based teaching of rural secondary science teachers through Authentic Research Experiences in oceanographic science and microbial ecology. The K20 network includes 96% of Oklahoma schools, providing a unique opportunity to impact STEM education in Oklahoma.

The results of this project will help develop a better understanding of DON cycling, the ecological context of creatine uptake activity, and identify both creatine-producing and consuming organisms in the marine environment. The importance of creatine cycling will be assessed via ¹⁵N tracer studies along the natural coastal-to-offshore productivity gradient observed in the North Atlantic. Tracer and molecular approaches will be used to investigate the importance of phytoplankton vs. bacteria in creatine uptake and, the taxonomic identities of creatine-utilizing bacteria will be interrogated via molecular, stable isotope probing (SIP), and RT-qPCR approaches.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1635369

[[table of contents](#) | [back to top](#)]