

Development (i.e. maturation) time measurements for *Acartia hudsonica* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA)

Website: <https://www.bco-dmo.org/dataset/924206>

Data Type: Other Field Results, experimental

Version: 1

Version Date: 2024-04-03

Project

» [Collaborative Research: Response of marine copepods to warming temperature and ocean acidification](#)
(Copepod Response to Warming Temp and OA)

Contributors	Affiliation	Role
Dam, Hans G.	University of Connecticut (UConn)	Principal Investigator
Baumann, Hannes	University of Connecticut (UConn)	Co-Principal Investigator
Finiguerra, Michael	University of Connecticut (UConn)	Co-Principal Investigator
Pespeni, Melissa	University of Vermont (UVM)	Co-Principal Investigator
Brennan, Reid	University of Vermont (UVM)	Scientist
deMayo, James	University of Connecticut (UConn)	Student
Park, Gihong	University of Connecticut (UConn)	Student
Norton, Lydia	University of Connecticut (UConn)	Technician
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

These data include development (i.e. maturation) time measurements for *Acartia hudsonica* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA) including a benign ambient condition temperature and CO₂ control (AM). These data were collected every second generation between F0 and F4 for all treatments and at F11 for AM and OWA. Data were calculated for naupliar development time (i.e. naupliar stage 1 to naupliar stage 6) and copepodite development time (i.e. copepodite stage 1 to adulthood). The experimental environmental conditions were: 1) Ambient control (AM): 13°C, 400 µatm CO₂, pH = 8.2; 2) Ocean Acidification (OA): 13°C, 1000 µatm CO₂, pH = 7.85; 3) Ocean Warming (OW): 15°C, 400 µatm CO₂, pH = 8.2; 4) Combined warming and acidification (OWA): 15°C, 1000 µatm CO₂, pH = 7.85.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Methods & Sampling](#)
 - [Data Processing Description](#)
 - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Location: University of Connecticut, Avery Point campus
Spatial Extent: Lat:41.3 Lon:-72
Temporal Extent: 2017-07-02 - 2018-08-15

Methods & Sampling

Three hundred copepods were collected in April of 2018 from eastern Long Island Sound, Connecticut, USA (41.3°N, 72.0°W) and raised for one year (~12 generations) (14 degrees Celsius (°C), 400 microatmospheres (µatm) CO2, 30 ‰ salinity, 12:12 hours light:dark) as stock cultures to limit maternal effects (Falconer, 1989, *Introduction to Quantitative Genetics*). Three resulting stock cultures with >2,000 individuals each were combined and then split evenly into three groups for each of the four treatments. Groups were acclimatized within a generation to temperature (15°C or 13°C, 1°C per day) and pCO2 (1000 µatm, 100 µatm per day, OA treatments only). Groups seeded the F0 individuals for 7-10 days yielding ~15,000 eggs per treatment. Resulting F0 eggs and nauplii were combined for each treatment, redistributed among three replicate cultures, and returned to their respective experimental conditions. The experimental environmental conditions were: 1) Ambient control (AM): 13°C, 400 µatm CO2, pH = 8.2; 2) Ocean Acidification (OA): 13°C, 1000 µatm CO2, pH = 7.85; 3) Ocean Warming (OW): 15°C, 400 µatm CO2, pH = 8.2; 4) Combined warming and acidification (OWA): 15°C, 1000 µatm CO2, pH = 7.85. Copepods were fed equal proportions of the live phytoplankters *Tetraselmis* sp., *Rhodomonas* sp., and *Thalassiosira weissflogii* every 48-72 hours to achieve food-replete conditions (≥600 micrograms (µg) Carbon per liter (L)) (Feinberg and Dam, 1998. *Marine Ecology Progress Series*), deliberately raised under ambient conditions to avoid confounding effects of possible food quality changes.

Development time was recorded as the number of days it took individuals to progress from N1 to C6 stage during the survival experiments. Individual development time values (N=34 - 187) were averaged across each treatment for each generation from F0-F4.

Data Processing Description

Data were processed and analyzed using R (v 4.0.2). Data analysis and visualization code can be accessed at: <https://zenodo.org/badge/latestdoi/505896789>.

BCO-DMO Processing Description

- Imported original file "Ahud_cdev_w_f11.txt" into the BCO-DMO system.
- Renamed fields to comply with BCO-DMO naming conventions.
- Saved final file as "924206_v1_a_hudsonica_development_time.csv".

[[table of contents](#) | [back to top](#)]

Data Files

File
924206_v1_a_hudsonica_development_time.csv (Comma Separated Values (.csv), 84.99 KB) MD5:25a43da5c9677aa28788e0fc31ddb919
Primary data file for dataset ID 924206, version 1

[[table of contents](#) | [back to top](#)]

Related Publications

R Core Team (2020). R: A language and environment for statistical computing. R v4.0.2. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
Software

deMayo, J. A., Brennan, R. S., Pespeni, M. H., Finiguerra, M., Norton, L., Park, G., Baumann, H., & Dam, H. G. (2023). Simultaneous warming and acidification limit population fitness and reveal phenotype costs for a marine copepod. *Proceedings of the Royal Society B: Biological Sciences*, 290(2006). <https://doi.org/10.1098/rspb.2023.1033>
Results

[[table of contents](#) | [back to top](#)]

Related Datasets

IsRelatedTo

Dam, H. G., Baumann, H., Finiguerra, M., Pespeni, M., Brennan, R. (2024) **Body size measurements collected for *Acartia hudsonica* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-04-03 doi:10.26008/1912/bco-dmo.924236.1 [[view at BCO-DMO](#)]

Relationship Description: These datasets result from the same set of experiments.

Dam, H. G., Baumann, H., Finiguerra, M., Pespeni, M., Brennan, R. (2024) **Egg production rate (EPR) and egg hatching success (HS) data for *Acartia tonsa* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-04-02 doi:10.26008/1912/bco-dmo.924126.1 [[view at BCO-DMO](#)]

Relationship Description: These datasets result from the same set of experiments.

Dam, H. G., Baumann, H., Finiguerra, M., Pespeni, M., Brennan, R. (2024) **Population fitness measurements collected for *Acartia hudsonica* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-03-29 doi:10.26008/1912/bco-dmo.923960.1 [[view at BCO-DMO](#)]

Relationship Description: These datasets result from the same set of experiments.

Dam, H. G., Baumann, H., Finiguerra, M., Pespeni, M., Brennan, R. (2024) **Population fitness measurements collected for *Acartia tonsa* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-03-28 doi:10.26008/1912/bco-dmo.923908.1 [[view at BCO-DMO](#)]

Relationship Description: These datasets result from the same set of experiments.

Dam, H. G., Baumann, H., Finiguerra, M., Pespeni, M., Brennan, R. (2024) **Survivorship measurements collected for *Acartia hudsonica* during multigenerational exposure to ocean warming (OW), ocean acidification (OA), and combined ocean warming and acidification (OWA).** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-04-01 doi:10.26008/1912/bco-dmo.924088.1 [[view at BCO-DMO](#)]

Relationship Description: These datasets result from the same set of experiments.

Jamesdemayo, & Dam-Lab. (2023). *dam-lab/hudsonica_transgenerational_MS: Data and code for manuscript: Limited copepod adaptation to combined warming and acidification reveals cost of producing adaptive phenotypes* (Version 3.3) [Computer software]. Zenodo. <https://doi.org/10.5281/ZENODO.6678641>
<https://doi.org/10.5281/zenodo.6678641>

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Treatment	The treatment experienced (1=AM, 2=OA, 3=OW, 4=OWA)	unitless
Temp	The temperature condition of the relevant treatment	degrees Celsius
pH	The pH condition of the relevant treatment	pH units
Rep	The biological culture replicate	unitless
Beak	The 25-mL beaker within a culture replicate	unitless
time	The day at which the beaker was measured	days post hatch
nx	The number of individuals at a particular time	number of individuals
lx	The proportion of surviving individuals at a particular time	number of individuals
Ndev	The number of individuals that matured from nauplii to copepodites on any given day	number of individuals
Cdev	The number of individuals that matured from copepodites to adulthood on any given day	number of individuals
F_Ratio	Ratio of females to total number of copepods in the population	females per total individuals
M_Ratio	Ratio of males to total number of copepods in the population	males per total individuals
Generation	The generation where the development time was measured	unitless
Generation_c	A numeric column for analyzing generation; equivalent to "Generation"	unitless

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Olympus SZH-ILLD Stereoscope
Generic Instrument Name	Microscope - Optical
Generic Instrument Description	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

[[table of contents](#) | [back to top](#)]

Project Information

Collaborative Research: Response of marine copepods to warming temperature and ocean acidification (Copepod Response to Warming Temp and OA)

Coverage: North western Atlantic ocean; Gulf of Maine, coastal and estuarine habitats

NSF Award Abstract:

Over time, our oceans are becoming both warmer and higher dissolved carbon dioxide. The latter condition is called ocean acidification. The consequences of these simultaneous changes for populations of marine organisms are not well understood. For this project, the investigators will conduct a series of laboratory

experiments to determine how two closely-related, common species of *Acartia* copepods will respond to the interactive effects of warming and acidification and also how well these species can adapt over multiple generations to changing ocean conditions. Since these copepods are key species in coastal food webs, results will have important implications for understanding and predicting how marine ecosystems may respond to future climate change. The investigators will share results from the research through traditional print media, case studies, and video mini lectures. The goal will be for educators of all levels to easily access material on climate change and ocean acidification to include in teaching curricula, in alignment with recommendations for universal design for learning. The project is a collaborative effort between an established professor at the University of Connecticut and an early-career female scientist at the University of Vermont. It will provide training and opportunities for collaborative, interdisciplinary research for two postdoctoral investigators, two graduate students and an undergraduate student.

The project's main goals are: 1) to test the simultaneous effects of temperature and carbon dioxide under current and future conditions on life history traits throughout the life cycle for two key copepod species, warm-adapted *Acartia tonsa* and cold-adapted *Acartia hudsonica*; 2) to test for adaptive capacity of both copepod species to a warmer and carbon-dioxide-enriched ocean; 3) to measure the genetic and maternally-induced changes across multiple generations of experimental selection in future conditions in both copepod species, and to identify the genes and pathways responding to selection. The investigators will use experiments encompassing current and projected temperature and carbon-dioxide conditions, will determine the roles of each variable and their interaction on traits that affect the fitness of both copepod species. They will also determine which life stages are most sensitive to individual or simultaneous stress conditions. Through multigenerational selection experiments, the investigators will identify and characterize the mechanisms of copepod evolutionary adaptation. Finally, they will measure genomic changes across the generations under all four experimental conditions to quantify the relative contributions of genetic and maternally-induced change in the physiological and life history traits of copepods in response to near-future climate conditions.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1559075
NSF Division of Ocean Sciences (NSF OCE)	OCE-1559180

[[table of contents](#) | [back to top](#)]