

# Pseudo-nitzschia spp. presence-absence and environmental data in Narragansett Bay in Rhode Island, USA and the Northeast U.S. Shelf (NES-LTER transect) from 2018-2023

Website: <https://www.bco-dmo.org/dataset/936856>

Data Type: Cruise Results, Other Field Results

Version: 1

Version Date: 2024-10-14

## Project

- » [Northeast U.S. Shelf Long Term Ecological Research site](#) (NES LTER)
- » [RII Track-1: Rhode Island Consortium for Coastal Ecology Assessment, Innovation, and Modeling](#) (C-AIM)
- » [Narragansett Bay Long-Term Plankton Time Series](#) (NBPTS)

## Program

- » [Long Term Ecological Research network](#) (LTER)

Contributors	Affiliation	Role
<a href="#">Jenkins, Bethany D.</a>	University of Rhode Island (URI)	Principal Investigator
<a href="#">Bertin, Matthew</a>	University of Rhode Island (URI)	Co-Principal Investigator
<a href="#">Kirk, Riley</a>	University of Rhode Island (URI)	Scientist
<a href="#">Rynearson, Tatiana A.</a>	University of Rhode Island (URI)	Scientist
<a href="#">Sterling, Alexa</a>	University of Rhode Island (URI)	Scientist
<a href="#">Church, Isabella</a>	University of Rhode Island (URI)	Student
<a href="#">Kim, Andrew</a>	University of Rhode Island (URI)	Student
<a href="#">Roche, Katherine M.</a>	University of Rhode Island (URI)	Student, Contact
<a href="#">Merchant, Lynne M.</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This dataset includes environmental measurements and presence-absence of *Pseudo-nitzschia* species, a harmful algal bloom diatom genus, associated with samples from various sites in Narragansett Bay, Rhode Island, including the Narragansett Bay Long Term Plankton Time Series site, and several stations along the Northeast U.S. Shelf Long Term Ecological Research program transect. These data correspond to an analysis of *Pseudo-nitzschia* species composition and domoic acid toxin production during winters and summers from 2018-2023 in Narragansett Bay and the Northeast U.S. Shelf, which was prepared for submission to Harmful Algae (Roche, et al.). This dataset includes sites information, particulate domoic acid concentration, *Pseudo-nitzschia* cell counts, temperature, salinity, nutrient concentrations, presence-absence of *Pseudo-nitzschia* species, and NCBI BioSample accessions.

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Methods & Sampling](#)
  - [Data Processing Description](#)
  - [BCO-DMO Processing Description](#)
- [Data Files](#)
- [Supplemental Files](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:41.57 E:-70.77 S:39.93 W:-71.42

**Temporal Extent:** 2018-01-31 - 2023-01-16

## Dataset Description

### Acknowledgement:

We acknowledge the NSF RI C-AIM EPSCoR Cooperative Agreement (OIA-1004057) for research support. Sequencing was performed at the University of Rhode Island Molecular Informatics Core supported by the Institutional Development Award (IDeA) Network for Biomedical Research Excellence from the National Institute of General Medical Sciences of the National Institutes of Health (P20GM103430).

## Methods & Sampling

Samples were selected from the NES-LTER transect and various time series sites in Narragansett Bay, Rhode Island (NB) during winter and summer periods from January 2018 through February 2023 to compare seasonal and regional patterns of *Pseudo-nitzschia* species composition and DA, as well as environmental drivers. NB and NES will henceforth be referred to as subregions of the larger Northeast U.S. Continental Shelf region, with NES specifically referring to the area spanned by the NES-LTER transect. Samples were collected on NES-LTER cruises (*R/V Endeavor*, *R/V Atlantis*) from 11 stations along a 150 km transect (n=77) each winter (January-February) and summer (July-August). Samples from three to four stations per cruise were used in this dataset spanning innershelf (L1), midshelf (L3, L4) and outershelf (L7, L8, L10) sections of the transect. The northernmost station, L1, is about 50 km from the mouth of NB. To collect plankton biomass for nucleic acid isolation, CTD rosette seawater from the surface and subsurface chlorophyll maximum (SCM) were passed via peristaltic pump over 25 mm 5 µm pore size filters (Sterlitech, Kent, WA, USA). Biomass filters were either flash frozen in liquid nitrogen (2018-2022) or placed in DNA/RNA shield (winter 2023; Zymo Research, Irvine, CA, USA) and stored in a -80°C freezer. The SCM depth varied as observed by *in situ* chlorophyll fluorescence, with a median depth of 28 m for summer and 19 m for winter samples. In cases where the SCM was not well defined due to water column mixing that typically took place in winter at nearshore stations, a sampling depth between 20 and 30 m was targeted.

In NB, surface seawater samples were collected from various sites in the East and West Passages including the Narragansett Bay Long-Term Plankton Time Series (NBPTS) site, Whale Rock (WR), Castle Hill Beach (CHB), East Passage (EP), and University of Rhode Island Graduate School of Oceanography (GSO) dock. Seawater was transported back to the laboratory and passed over 25 mm 5 µm pore size filters (Sterlitech, Kent, WA, USA) using a peristaltic pump before flash freezing in liquid nitrogen and storage at -80°C. To fill in several missing dates from this time series, six samples collected separately in the NBPTS (<https://web.uri.edu/gso/research/plankton/>) sampling program were used. These samples differed in collection methodology only by the filter pore size used (0.22 µm, Express Plus, Millipore Sigma) and vacuum as opposed to peristaltic filtration.

DNA was extracted from most NB and NES samples (n=219) using a modified version of the DNeasy Plant Kit (Qiagen, Germantown, MD, USA) that included a 1-minute bead

beating step (0.1 mm and 0.5 mm Zirconia/Silica beads, BioSpec Products, Bartlesville, OK, USA) and two part elution into a total of 45 µL Buffer AE. Similarly, the six NBPTS samples were extracted using a modified version of the DNeasy Blood & Tissue kit (Qiagen, Germantown, MD, USA) with a 1-minute bead beating step and final elution into 50 µL Buffer AE. Some NES samples (n=18) were extracted using the Quick-DNA/RNA Miniprep Plus Kit (Zymo Research, Irvine, CA, USA) with a 1-minute bead beating step (0.4 mm Zirconium Beads, OPS Diagnostics, Lebanon, NJ, USA) and final elution into 50 µL nuclease-free water. DNA from each sample was amplified with a primer set that targets the eukaryotic internal transcribed spacer region 1 (ITS1) and effectively distinguishes *Pseudo-nitzschia* species (White et al., 1990; Sterling et al., 2022). Briefly, DNA was diluted to 1-4 ng/µL and 2 µL of template was added to 25 µL PCR reactions with Phusion Hot Start High-Fidelity Master Mix (Thermo Fisher Scientific Inc., Waltham, MA, USA) and HPLC-purified forward and reverse primers at 0.5 µM concentration with Illumina MiSeq adapters (Integrated DNA Technologies, Coralville, IA, USA). A stepwise thermocycle was used as described in Sterling et al. (2022).

DNA amplicons were sequenced at the Rhode Island-INBRE Molecular Informatics Core on the Illumina MiSeq platform (Illumina, Inc., San Diego, CA, USA). There, libraries were prepared by cleaning ITS1 PCR products with KAPA pure beads (KAPA Biosystems, Woburn, MA, USA) and attaching sequencing indices and adapters using PCR. This amplification was performed with the Illumina Nextera XT Index Kit (Illumina, San Diego, CA, USA) and Phusion High Fidelity Master Mix, followed by a second round of cleaning with KAPA pure beads and visualization with gel electrophoresis. The quality of select samples was assessed on a Bioanalyzer DNA1000 chip (Agilent Technologies, Santa Clara, CA, USA) and all samples were quantified on a Qubit fluorometer (Invitrogen, Carlsbad, CA, USA). The final library was pooled, quantified with qPCR on a LightCycler480 (Roche, Pleasanton, CA, USA) using a KAPA Biosystems Illumina Kit (KAPA Biosystems, Woburn, MA, USA), and sequenced on the Illumina MiSeq using v3 chemistry and 2x250 paired-end reads. Samples were sequenced across five separate MiSeq runs using identical methods and negative controls.

Various biological, chemical, and physical data were collected during each sampling event. In NB, surface temperature and salinity were measured using multiparameter sondes (6920 V2 for samples collected at NBPTS and WR; ProDSS for samples collected at CHB and GSO dock; YSI, Yellow Springs, Ohio, USA). During NES sampling, temperature and salinity were measured using two SBE911 CTD sensors (Sea Bird Electronics, Bellevue, WA, USA) and the mean of the two measurements was used in the final analysis. Dissolved macronutrient samples were collected by freezing 0.2 µm seawater filtrate at -20°C. NB site nutrients were analyzed at the University of Rhode Island Marine Science Research Facility (URI MSRF, Narragansett, RI, USA) on a QuickChem 8500 (Lachat, Milwaukee, WI, USA) while NES samples were measured at the Woods Hole Oceanographic Institution's Nutrient Analytical Facility (Woods Hole, MA, USA) on a four-channel segmented flow AA3 HR Autoanalyzer (SEAL Analytical, Mequon, WI, USA). Both instruments measured nitrite + nitrate, ammonium, silicate, and phosphate. Nitrite + nitrate and ammonium values were summed and used in the analysis as dissolved inorganic nitrogen (DIN). Any measurements below each instrument's limit of detection for each nutrient type were replaced with zero.

Biomass for particle-associated DA analysis was collected, extracted, and analyzed via liquid chromatography with tandem mass spectrometry (LC-MS/MS) with multiple reaction monitoring. All samples were chromatographically separated in an identical fashion to Sterling et al. (2022), and the majority of samples (n=167) were analyzed using a 4500 QTRAP mass spectrometer (SCIEX, Framingham, MA, USA). A subset of samples (n=42) were measured using a 1290 Infinity II UHPLC system coupled to a 6470 Triple Quadrupole mass spectrometer (Agilent Technologies, Santa Clara, CA, USA). The peak of DA eluted at 10.41 min. Analysis was carried out in positive mode, and three transitions from the protonated DA molecule were used and optimized for quantification:  $m/z$  312 → 266,  $m/z$  312 → 161, and  $m/z$  312 → 105 determined by MassHunter Optimizer (Agilent, Santa Clara, CA, USA) including the optimized fragmentor (112), collision energy (20, 28, 48), and cell accelerator voltage (4) settings. The  $m/z$  312 → 266 transition was used for quantification following acquisition from both mass spectrometry instruments. Particle-associated DA was quantified to ng particulate DA L<sup>-1</sup> of filtered seawater using an external calibration curve created with pure DA standards of increasing concentrations included in each analysis (DA Certified Reference Material, National Research Council Canada, Halifax, Nova Scotia).

*Pseudo-nitzschia* spp. abundance was quantified using light microscopy cell counts of live (NBPTS site) and preserved (all other sites) samples. For preserved samples, acidic Lugol's solution was added to whole seawater for a final concentration of 1% Lugol's and stored at 4°C until enumeration. A Sedgewick-Rafter counting chamber (Science First/Wildco, Yulee, FL, USA) and a BX40 light microscope (Olympus, Tokyo, Japan) were used to identify and enumerate cells at the genus level, since many *Pseudo-nitzschia* species are morphologically cryptic under light microscopy (Bates et al., 2018).

#### Sampling Locations

Narragansett Bay sites:

Narragansett Bay Long Term Plankton Time Series (41.57 N -71.39 W)

Whale Rock (41.43 N -71.42 W)

East Passage (41.45 N -71.38 W)

Castle Hill Beach (41.46 N -71.36 W)

Graduate School of Oceanography dock (41.49 N -71.42 W)

NES stations:

L1 (41.20 N -70.88 W), L3 (40.86 N -70.88 W), L4 (40.70 N -70.88 W), L7 (40.23 N -70.88 W), L8 (40.14 N -70.77 W), L10 (39.93 N -70.88 W)

#### Data Processing Description

Raw sequencing read quality was assessed using FastQC and MultiQC (v.0.11.9, v1.11) before and after primer and adapter trimming in Cutadapt (v3.2). The Divisive Amplicon Denoising Algorithm (DADA2) was used in R to estimate sequencing error and identify distinct amplicon sequence variants (ASVs) (v1.20.0). DADA2 was run separately for each sequencing run because it is designed to account for run-specific error. Taxonomy was assigned to ASVs from all sequencing runs using a dual approach to maximize the number of ASVs identified to the species level and enhance confidence. First, the scikit-learn naïve Bayes machine learning classifier in QIIME2 (v2022.11) and a curated reference database were used to assign taxonomy with a confidence threshold of 0.8. This curated database from Roche et al. (2022) included 302 unique *Pseudo-nitzschia* spp. ITS1 sequences from the National Center for Biotechnology Information (NCBI) GenBank with 51 different species represented (retrieved June 1, 2021). Next, to ensure that relevant *Pseudo-nitzschia* spp. ITS1 taxonomy was not omitted from the curated database, a megablast search was performed using the entire BLAST nt database. Additional ASVs classified by megablast were retained if there was >99% identity and >99% query cover to NCBI *Pseudo-nitzschia* species. If QIIME2 and megablast taxonomic assignment did not match for a particular ASV, no species-level taxonomy was assigned. Samples containing no reads identified to the *Pseudo-nitzschia* species level were removed from the analysis (n=15). To avoid potentially falsely detected taxa, ASVs classified as a species that appeared in only one sample across the dataset were discarded (n=6).

#### BCO-DMO Processing Description

1) Processed the submitted file named DATASET02\_env\_metadata\_species\_presence\_v2.csv, which contains environmental data and indications of the presence and absence of species, with the BCO-DMO tool Laminar.

Renamed parameters according to BCO-DMO naming conventions. Replaced all periods in the parameter names with underscores and removed units from names.

Converted the format of the sample date parameter 'Sample\_Date' from %m/%d/%y to and ISO 8601 standard format of %Y-%m-%d.

Saved the modified dataset to the file 936856\_v1\_pseudo\_nitzschia\_environmental.csv

2) Processed the submitted file named DATASET02\_env\_metadata\_species\_presence\_v2.csv with Laminar to create an unpivoted format.

Renamed parameters according to BCO-DMO naming conventions. Replaced all periods in the parameter names with underscores and removed units from names.

Converted the format of the sample date parameter 'Sample\_Date' from %m/%d/%y to and ISO 8601 standard format of %Y-%m-%d.

Unpivoted the dataset on the species parameters to produce two columns named species and present. The column 'species' lists all the species parameter names and the column

'present' is a flag representing 1 if a species is present and 0 if a species is absent.

The values in the species column were modified to replace underscores with spaces, add a period after var, and add a hyphen between Pseudo and nitzschia to get Pseudo-nitzschia.

Saved the modified dataset to the file unpivoted\_pseudo\_nitzschia\_environmental.csv.

3) Renamed the submitted file Supp\_DATASET\_PNsSpecies\_presence\_absence.csv, which contains indications of the presence and absence of species, to pseudo\_nitzschia\_presence\_absence.csv.

4) Created a taxonomy table using the species names in the dataset and getting taxonomy values using the World Register of Marine Species (WoRMS) website and saved it to the file species\_taxonomy.csv.

[ [table of contents](#) | [back to top](#) ]

## Data Files

File
<b>Environmental data, Pseudo-nitzschia species presence-absence, and NCBI accessions</b> filename: 936856_v1_pseudo_nitzschia_environmental.csv (Comma Separated Values (.csv), 51.48 KB) MD5:c053875db723cb3e56808b042d99a975 Primary data file for dataset ID 936856, version 1 This file contains metadata, environmental data, Pseudo-nitzschia species presence-absence, and NCBI accession numbers for sequencing data

[ [table of contents](#) | [back to top](#) ]

## Supplemental Files

File
<b>Parameter descriptions for the unpivoted file</b> filename: parameter_descriptions_for_unpivoted_file.csv (Comma Separated Values (.csv), 3.14 KB) MD5:4ad5b733f274a44b1480bd62b8ce613 Parameter descriptions for parameters in the supplemental data file "Unpivoted Pseudo-nitzschia species environmental with presence and absence"
<b>Pseudo-nitzschia species presence and absence</b> filename: pseudo_nitzschia_presence_absence.csv (Comma Separated Values (.csv), 10.96 KB) MD5:b6a687900f452997714cf96a817c225 Presence and absence of Pseudo-nitzschia species Species column and Library ID columns ranging from IC01 - IC40, AS304 - AS496, KR133 - KR156, IC109 - IC237, KR140 - KR146, AS328 - AS436, KR702 - KR735 The Library ID column names are the library_ID values found in the primary data file. Library ID parameter description: Sequencing identifying number that associates environmental data with sequencing data matrix
<b>Species taxonomy</b> filename: species_taxonomy.csv (Comma Separated Values (.csv), 3.75 KB) MD5:74f167cc0e1a756b9a9f900c6173b02 Species taxonomy from the World Register of Marine Species (WoRMS) with columns: ScientificName, AphiaID, LSID, Kingdom, Phylum, Class, Order, Family, Genus, Species Parameter descriptions ScientificName: Genus and Species name AphiaID: Unique taxonomic identifier at the World Register of Marine Species (WoRMS: marinespecies.org) LSID: The Life Sciences Identifier (LSID) is an Interoperable Informatics Infrastructure Consortium (I3C) and OMG Life Sciences Research (LSR) Uniform Resource Name (URN) specification in progress.
<b>Unpivoted Pseudo-nitzschia species environmental with presence and absence</b> filename: unpivoted_pseudo_nitzschia_environmental.csv (Comma Separated Values (.csv), 1,007.34 KB) MD5:b4371eeed5180dd45ec6e3b04bcc7a75 Unpivoted version of the primary data file For parameter descriptions, see the supplemental file "parameter_descriptions_for_unpivoted_file.csv".

[ [table of contents](#) | [back to top](#) ]

## Related Publications

Andrews S. (2010). FastQC: a quality control tool for high throughput sequence data. Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>  
*Software*

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... Asnicar, F. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, 37(8), 852-857. doi:[10.1038/s41587-019-0209-9](https://doi.org/10.1038/s41587-019-0209-9)  
*Software*

Ewels, P., Magnusson, M., Lundin, S., & Källér, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19), 3047-3048. doi:[10.1093/bioinformatics/btw354](https://doi.org/10.1093/bioinformatics/btw354)  
*Software*

FastQC (2015), FastQC [Online]. Available online at: <https://qubeshub.org/resources/fastqc>.  
*Software*

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 17(1), 10. doi:[10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200)  
*Software*

R Core Team (2023). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>  
*Software*

Roche, K. M., Sterling, A. R., Ryneerson, T. A., Bertin, M. J., & Jenkins, B. D. (2022). A Decade of Time Series Sampling Reveals Thermal Variation and Shifts in Pseudo-nitzschia Species Composition That Contribute to Harmful Algal Blooms in an Eastern US Estuary. *Frontiers in Marine Science*, 9. <https://doi.org/10.3389/fmars.2022.889840>  
*Methods*

Roche, K.M., Church, I.N., Sterling, A.R., Ryneerson, T.A., Bertin, M.J., Kim, A.M., Kirk, R.D., Jenkins, B.D. (2024). Connectivity of toxigenic Pseudo-nitzschia species assemblages between the Northeast U.S. continental shelf and an adjacent estuary. Manuscript submitted for publication.  
*Results*

Sterling, A. R., Kirk, R. D., Bertin, M. J., Ryneerson, T. A., Borkman, D. G., Caponi, M. C., Carney, J., Hubbard, K. A., King, M. A., Maranda, L., McDermith, E. J., Santos, N. R., Strock, J. P., Tully, E. M., Vaverka, S. B., Wilson, P. D., & Jenkins, B. D. (2022). Emerging harmful algal blooms caused by distinct seasonal assemblages of a toxic diatom. *Limnology and Oceanography*, 67(11), 2341-2359. Portico. <https://doi.org/10.1002/lno.12189>  
*Methods*

White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). AMPLIFICATION AND DIRECT SEQUENCING OF FUNGAL RIBOSOMAL RNA GENES FOR PHYLOGENETICS. *PCR Protocols*, 315-322. <https://doi.org/10.1016/b978-0-12-372180-8.50042-1>  
*Methods*

[ [table of contents](#) | [back to top](#) ]

## Related Datasets

### IsRelatedTo

Roche, K. M., Church, I., Sterling, A., Ryneerson, T. A., Bertin, M., Kim, A., Kirk, R., Jenkins, B. D. (2024) **Amplicon sequence variants (ASVs) and taxonomy of Pseudo-nitzschia spp. from Narragansett Bay in Rhode Island, USA and the Northeast U.S. Shelf (NES-LTER transect) from 2018-2023.** Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2024-10-11 doi:10.26008/1912/bco-dmo.936849.1 [[view at BCO-DMO](#)]

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
library_ID	Sequencing identifying number that associates environmental data with sequencing data matrix.	unitless
Sample_Date	Sample collection date	unitless
Site	Sample collection site name	unitless
Site_Abbreviation	Sample collection site name abbreviated	unitless
Lat	Latitude of sampling site (N = positive)	decimal degrees
Lon	Longitude of sampling site (W = negative)	decimal degrees
Depth_category	Qualitative depth category used in analysis to indicate sampling depth. Surface means that the sample was collected within the upper 5 m of the water column and SCM stands for subsurface chlorophyll maximum which was determined by chlorophyll fluorescence.	unitless
Depth	Depth of sample collection	meters
Season	Season of sample collection used in analysis	unitless
LTER_cruise	NES LTER cruise deployment name. NA = not applicable	unitless
CTD_cast	Number of CTD cast on NES LTER cruise. One sample indicates "flow" because it was collected from the ship flow-through system instead of a CTD cast. One sample indicates "bucket" because it was collected using a bucket and rope from surface seawater. NA = not applicable	unitless
Niskin	Number of niskin bottle from CTD casts on NES LTER cruise. One sample indicates "flow" because it was collected from the ship flow-through system instead of a CTD cast. One sample indicates "bucket" because it was collected using a bucket and rope from surface seawater. NA = not applicable	unitless
pDA	Particulate domoic acid concentration measured by LC-MS/MS of > 5um filters	nanograms/liter (ng/L)
Pseudonitzschia	Number of individual cells identified as belonging to the diatom genus Pseudo-nitzschia [NCBI:txid41953] calculated in one liter (L) of surface seawater	cells/liter
Temp	Seawater temperature	Degrees Celsius (°C)
Salinity	Seawater salinity	Practical Salinity Units (PSU)
Phosphate	Concentration of dissolved inorganic phosphate in seawater sample	microMolar (uM)
Silicate	Concentration of dissolved inorganic silicate in seawater sample	microMolar (uM)
Nitrate_and_Nitrite	Concentration of dissolved nitrate and nitrite in seawater sample	microMolar (uM)
Nitrite	Concentration of dissolved nitrite in seawater sample	microMolar (uM)
Ammonium	Concentration of dissolved ammonium in seawater sample	microMolar (uM)
Nitrate	Concentration of dissolved nitrate in seawater sample calculated by subtracting the concentration of nitrite from the nitrate plus nitrite measurement	microMolar (uM)
DIN	Concentration of dissolved inorganic nitrogen in seawater sample calculated by summing the nitrate plus nitrite and ammonium measurements	microMolar (uM)

Pseudo_nitzschia_americana	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_australis	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_caciantha	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_calliantha	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_cuspidata	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_delicatissima	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_fraudulenta	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_galaxiae	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_hasleana	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_inflatula	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_mannii	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_multiseries	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_multistriata	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_plurisetata	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_pungens	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_pungens_var_aveirensis	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_pungens_var_cingulata	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_qiana	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_sabit	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_subpacificata	presence (1) or absence (0) of the species in sample	unitless
Pseudo_nitzschia_turgidula	presence (1) or absence (0) of the species in sample	unitless
NCBI_BioSample	NCBI BioSample accession number	unitless
NCBI_BioProject	NCBI BioProject number	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	Illumina MiSeq Platform
<b>Generic Instrument Name</b>	Automated DNA Sequencer
<b>Dataset-specific Description</b>	Manufactured by Illumina, Inc., San Diego, CA, USA
<b>Generic Instrument Description</b>	General term for a laboratory instrument used for deciphering the order of bases in a strand of DNA. Sanger sequencers detect fluorescence from different dyes that are used to identify the A, C, G, and T extension reactions. Contemporary or Pyrosequencer methods are based on detecting the activity of DNA polymerase (a DNA synthesizing enzyme) with another chemoluminescent enzyme. Essentially, the method allows sequencing of a single strand of DNA by synthesizing the complementary strand along it, one base pair at a time, and detecting which base was actually added at each step.

<b>Dataset-specific Instrument Name</b>	SBE911 CTD sensors
<b>Generic Instrument Name</b>	CTD Sea-Bird 911
<b>Dataset-specific Description</b>	Manufactured by Sea Bird Electronics, Bellevue, WA, USA
<b>Generic Instrument Description</b>	The Sea-Bird SBE 911 is a type of CTD instrument package. The SBE 911 includes the SBE 9 Underwater Unit and the SBE 11 Deck Unit (for real-time readout using conductive wire) for deployment from a vessel. The combination of the SBE 9 and SBE 11 is called a SBE 911. The SBE 9 uses Sea-Bird's standard modular temperature and conductivity sensors (SBE 3 and SBE 4). The SBE 9 CTD can be configured with auxiliary sensors to measure other parameters including dissolved oxygen, pH, turbidity, fluorescence, light (PAR), light transmission, etc.). More information from Sea-Bird Electronics.

<b>Dataset-specific Instrument Name</b>	Lachat QuickChem 8500
<b>Generic Instrument Name</b>	Lachat QuikChem 8500 flow injection analysis system
<b>Dataset-specific Description</b>	Manufactured by Hach, Loveland, CO, USA
<b>Generic Instrument Description</b>	The Lachat QuikChem 8500 Series 2 Flow Injection Analysis System features high sample throughput and simple, but rapid, method changeover. The QuikChem 8500 Series 2 system maximises productivity in determining ionic species in a variety of sample types, from sub-ppb to percent concentrations. Analysis takes 20 to 60 seconds, with a sample throughput of 60 to 120 samples per hour.

<b>Dataset-specific Instrument Name</b>	SCIEX 4500 Qtrap mass spectrometer
<b>Generic Instrument Name</b>	Mass Spectrometer
<b>Dataset-specific Description</b>	Manufactured by Sciex, Framingham, MA, USA
<b>Generic Instrument Description</b>	General term for instruments used to measure the mass-to-charge ratio of ions; generally used to find the composition of a sample by generating a mass spectrum representing the masses of sample components.

<b>Dataset-specific Instrument Name</b>	6470 Triple Quadrupole mass spectrometer
<b>Generic Instrument Name</b>	Mass Spectrometer
<b>Dataset-specific Description</b>	Manufactured by Agilent Technologies, Santa Clara, CA, USA
<b>Generic Instrument Description</b>	General term for instruments used to measure the mass-to-charge ratio of ions; generally used to find the composition of a sample by generating a mass spectrum representing the masses of sample components.

<b>Dataset-specific Instrument Name</b>	BX40 light microscope
<b>Generic Instrument Name</b>	Microscope - Optical
<b>Dataset-specific Description</b>	Manufactured by Olympus, Tokyo, Japan
<b>Generic Instrument Description</b>	Instruments that generate enlarged images of samples using the phenomena of reflection and absorption of visible light. Includes conventional and inverted instruments. Also called a "light microscope".

<b>Dataset-specific Instrument Name</b>	AA3 HR Autoanalyzer
<b>Generic Instrument Name</b>	Seal Analytical AutoAnalyser 3HR
<b>Dataset-specific Description</b>	Manufactured by SEAL Analytical, Mequon, WI, USA
<b>Generic Instrument Description</b>	A fully automated Segmented Flow Analysis (SFA) system, ideal for water and seawater analysis. It comprises a modular system which integrates an autosampler, peristaltic pump, chemistry manifold and detector. The sample and reagents are pumped continuously through the chemistry manifold, and air bubbles are introduced at regular intervals forming reaction segments which are mixed using glass coils. The AA3 uses segmented flow analysis principles to reduce inter-sample dispersion, and can analyse up to 100 samples per hour using stable LED light sources.

<b>Dataset-specific Instrument Name</b>	Sedgewick Rafter Counting Chamber
<b>Generic Instrument Name</b>	Sedgewick Rafter Counting Chamber
<b>Dataset-specific Description</b>	Manufactured by Science First/Wildco, Yulee, FL, USA
<b>Generic Instrument Description</b>	Sedgewick Rafter Counting Chambers are transparent slides widely water analysis, culture inspection, and for other applications where particles per unit volume in fluid must be determined. The slide has a base that is ruled in one-thousand 1-millimeter squares. When a liquid is held in the cell by a coverglass, the grid subdivides 1 milliliter of liquid into 1 microliter volume.

<b>Dataset-specific Instrument Name</b>	Eppendorf Mastercycler EP Thermal Cyclers Series
<b>Generic Instrument Name</b>	Thermal Cyclers
<b>Generic Instrument Description</b>	A thermal cycler or "thermocycler" is a general term for a type of laboratory apparatus, commonly used for performing polymerase chain reaction (PCR), that is capable of repeatedly altering and maintaining specific temperatures for defined periods of time. The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. They can also be used to facilitate other temperature-sensitive reactions, including restriction enzyme digestion or rapid diagnostics. (adapted from <a href="http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html">http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html</a> )

<b>Dataset-specific Instrument Name</b>	1290 Infinity II UHPLC system
<b>Generic Instrument Name</b>	Ultra high-performance liquid chromatography
<b>Dataset-specific Description</b>	Manufactured by Agilent Technologies, Santa Clara, CA, USA
<b>Generic Instrument Description</b>	Ultra high-performance liquid chromatography: Column chromatography where the mobile phase is a liquid, the stationary phase consists of very small (< 2 microm) particles and the inlet pressure is relatively high.

<b>Dataset-specific Instrument Name</b>	YSI ProDSS multiparameter meter
<b>Generic Instrument Name</b>	YSI Professional Plus Multi-Parameter Probe
<b>Dataset-specific Description</b>	Manufactured by YSI Inc. / Xylem Inc., Yellow Springs, OH, USA
<b>Generic Instrument Description</b>	The YSI Professional Plus handheld multiparameter meter provides for the measurement of a variety of combinations for dissolved oxygen, conductivity, specific conductance, salinity, resistivity, total dissolved solids (TDS), pH, ORP, pH/ORP combination, ammonium (ammonia), nitrate, chloride and temperature. More information from the manufacturer.

[ [table of contents](#) | [back to top](#) ]

## Deployments

### EN608

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/848016">https://www.bco-dmo.org/deployment/848016</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2018-01-31
<b>End Date</b>	2018-02-06
<b>Description</b>	C-AIM project

### EN617

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/848018">https://www.bco-dmo.org/deployment/848018</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2018-07-20
<b>End Date</b>	2018-07-25

### EN627

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/848056">https://www.bco-dmo.org/deployment/848056</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2019-02-01
<b>End Date</b>	2019-02-06

### EN644

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/848020">https://www.bco-dmo.org/deployment/848020</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2019-08-20
<b>End Date</b>	2019-08-25

### EN649

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940001">https://www.bco-dmo.org/deployment/940001</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2020-02-01
<b>End Date</b>	2020-02-06
<b>Description</b>	Project: NES-LTER # 4

### EN655

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940004">https://www.bco-dmo.org/deployment/940004</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2020-07-25
<b>End Date</b>	2020-07-28
<b>Description</b>	Project: NES-LTER

#### EN661

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940007">https://www.bco-dmo.org/deployment/940007</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2021-02-03
<b>End Date</b>	2021-02-08
<b>Description</b>	Project: NES-LTER transect #8

#### EN668

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940010">https://www.bco-dmo.org/deployment/940010</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2021-07-16
<b>End Date</b>	2021-07-21
<b>Description</b>	Project: NES-LTER transect #9

#### AT46

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940019">https://www.bco-dmo.org/deployment/940019</a>
<b>Platform</b>	R/V Atlantis
<b>Start Date</b>	2022-02-16
<b>End Date</b>	2022-02-21
<b>Description</b>	Project: LTER: Linking Pelagic Community Structure with Ecosystem Dynamics and Production Regimes on the Changing Northeast US Shelf

#### EN687

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940013">https://www.bco-dmo.org/deployment/940013</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2022-07-29
<b>End Date</b>	2022-08-03
<b>Description</b>	Project: NES - LTER Summer 2022

#### EN695

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/940016">https://www.bco-dmo.org/deployment/940016</a>
<b>Platform</b>	R/V Endeavor
<b>Start Date</b>	2023-01-11
<b>End Date</b>	2023-01-16
<b>Description</b>	Project: NES-LTER 2023-01

[ [table of contents](#) | [back to top](#) ]

## Project Information

### Northeast U.S. Shelf Long Term Ecological Research site (NES LTER)

**Website:** <https://nes-lter.who.edu/>

**Coverage:** Northeast U.S. Continental Shelf Large Marine Ecosystem: 35.2019 to 46.0906 latitude, -77.3492 to -63.3608 longitude

### Continuing Award OCE-2322676

Sep 2023 to Aug 2028 (estimated)

**LTER: Scales of Variability in Ecosystem Dynamics and Production on the Changing Northeast U.S. Shelf (NES II)**

**NSF Award Abstract:**

The Northeast U.S. Shelf (NES) is the region of the Northwest Atlantic Ocean that overlies the continental shelf from North Carolina to Maine. The NES has a long history of intense human utilization and provides an array of ecosystem services including shipping, recreation, conservation, and energy development. The NES also comprises a seasonally dynamic and productive ecosystem, supporting renowned fisheries, whose integrity is critical to the health of the Northeast U.S. economy. The NES ecosystem's productivity is fueled by planktonic organisms that interact with each other in complex food webs whose structure depends on environmental conditions (e.g., temperature, light, and nutrient levels). These conditions are rapidly changing because of climate-change-related warming and human utilization. For example, the NES is seeing the largest development of coastal wind farms in the U.S. to date. Phase II of the Northeast U.S. Shelf Long-Term Ecological Research program (NES-LTER II) advances our ability to predict how anthropogenic impacts will affect the dynamics of the shelf's planktonic food webs and their ability to support the productivity of higher trophic levels, from fish to whales and humans. Because the NES is subject to long-term challenges that will impact many people, the project emphasizes an active education component for helping to train the next generation of marine scientists and outreach activities to increase public understanding of marine science and technology. The project team conducts education and outreach via three main components: (1) training and mentoring for early career researchers from undergraduates to postdoctoral researchers in LTER research; (2) an LTER Schoolyard program that engages middle and high school teachers and students; and (3) public outreach through targeted events, the project website, and social media channels.

Patterns of ecosystem change over seasons to decades have been documented in the NES, but the key mechanisms linking changes in the physical environment, planktonic food webs, and higher trophic levels remain poorly understood. As a result, predictive capability is limited and management strategies are largely reactive. To address these needs, NES II is targeting a mechanistic understanding of how food web structure and function responds to environmental conditions, natural variability and human induced changes. NES II combines observations that provide regional-scale context, process cruises along a high gradient cross-shelf transect, high-frequency time series at an inner-shelf location, coupled biological-physical food web models, and targeted population models. In addition, the research team is investigating how community structure and trophic transfer are



impacted by disturbances including (i) the increasing prevalence of heat waves, (ii) intrusions of offshore water associated with increasing instability in the Gulf Stream, and (iii) offshore wind farms now under construction on the NES. The long-term research plan is guided by the overarching science question: "How is climate change impacting the pelagic NES ecosystem and, in particular, affecting the relationship between compositional (e.g., species diversity and size structure) and aggregate (e.g., rates of primary production, and transfer of energy to higher trophic levels) variability?" The investigators are assessing the extent to which the NES ecosystem possesses a biodiversity reservoir that is resilient to dramatic changes in the environment and that will allow the ecosystem to maintain overall productivity.

#### **Prior Award**

**Sep 2017 to Feb 2024**

#### **LTERR: Linking Pelagic Community Structure with Ecosystem Dynamics and Production Regimes on the Changing Northeast US Shelf**

Summary information including abstract, PIs, and other award details are included in the Funding History PDF in the Files section below.

#### **Additional Information:**

The NES-LTERR project includes collaboration with the National Marine Fisheries Service / Northeast Fisheries Science Center [NMFS/NEFSC] in particular for sharing data related to Project EcoMon Zooplankton <https://www.bco-dmo.org/project/2106>.

#### **This project is supported by continuing grants with slight name variations:**

- LTER: Linking Pelagic Community Structure with Ecosystem Dynamics and Production Regimes on the Changing Northeast US Shelf
- LTER: Scales of Variability in Ecosystem Dynamics and Production on the Changing Northeast U.S. Shelf (NES II)

#### **RII Track-1: Rhode Island Consortium for Coastal Ecology Assessment, Innovation, and Modeling (C-AIM)**

**Coverage:** Narragansett Bay, Rhode Island

#### **NSF Award Abstract:**

##### **Non-technical Description**

The University of Rhode Island (URI) will establish the Consortium for Coastal Ecology Assessment, Innovation, and Modeling (C-AIM) to coordinate research, education, and workforce development across Rhode Island (RI) in coastal marine science and ecology. C-AIM addresses fundamental research questions using observations, computational methods, and technology development applied to Narragansett Bay (NB), the largest estuary in New England and home to important ecosystem services including fisheries, recreation, and tourism. The research will improve understanding of the microorganisms in NB, develop new models to predict pollution and harmful algal bloom events in NB, build new sensors for nutrients and pollutants, and provide data and tools for stakeholders in the state. Observational capabilities will be coordinated in an open platform for researchers across RI; it will provide real-time physical, chemical, and biological observations including live streaming to mobile devices. C-AIM will also establish the RI STEAM (STEM + Art) Imaging Consortium to foster collaboration between artists, designers, engineers, and scientists. Research internships will be offered to undergraduate students throughout the state and seed funding for research projects will be competitively awarded to Primarily Undergraduate Institution partners.

##### **Technical Description**

C-AIM will employ observations and modeling to assess interactions between organisms and ecosystem function in NB and investigate ecological responses to environmental events, such as hypoxia and algal blooms. Observations of the circulation, biogeochemistry, and ecosystem will be made using existing and new instrument platforms. The Bay Observatory ? a network of observational platforms around NB - will be networked to trigger enhanced water sampling and sensing during specific environmental events, such as hypoxic conditions or phytoplankton blooms. Biogeochemical, ecological, and coastal circulation models will be integrated and coupled to focus on eutrophication and pollutant loading. Data and models will be integrated on multiple scales, from individual organisms and trophic interactions to food-web responses, and from turbulence to the regional ocean circulation. New sensing technologies for nutrients and pollutants will be developed, including affordable, micro-fluidic (Lab-on-a-Chip) devices with antifouling capabilities. The results will be synthesized and communicated to stakeholders.

#### **Narragansett Bay Long-Term Plankton Time Series (NBPTS)**

**Website:** <https://web.uri.edu/gso/research/plankton/>

The Narragansett Bay Long-Term Plankton Time Series is one of the world's longest-running plankton surveys. Beginning in 1957, weekly samples have been collected to assess the phytoplankton community and characterize the physical parameters of Narragansett Bay.

Samples are collected once per week -regardless of tidal stage- for temperature, salinity, turbidity, size-fractionated chlorophyll a and nutrients. Microplankton community composition (size range >10µm, both species identification and abundance) is determined using a light microscope to quantify live samples. The species list for the >10µm size fraction includes 246 different species or species complexes of protists. Samples are also collected for the determination of copepod and ctenophore concentrations.

Funding for the time series has come from the University of Rhode Island since 1999. Ship time is frequently provided by the U.S. Department of Fish and Wildlife.

#### **This Time Series is related to the following projects at BCO-DMO:**

- Connecting local, regional and global scales of gene flow in planktonic marine diatoms (<https://www.bco-dmo.org/project/511708>)
- Dimensions: Collaborative Research: Genetic, functional and phylogenetic diversity determines marine phytoplankton community responses to changing temperature and nutrients (<https://www.bco-dmo.org/project/712787>)
- LTER: Linking Pelagic Community Structure with Ecosystem Dynamics and Production Regimes on the Changing Northeast US Shelf (<https://www.bco-dmo.org/project/747769>)
- Quantifying Temperature Dependence In Growth & Grazing Rates of Planktonic Herbivores (<https://www.bco-dmo.org/project/739232>)
- RII Track-1: Rhode Island Consortium for Coastal Ecology Assessment, Innovation, and Modeling (<https://www.bco-dmo.org/project/836631>)

[ [table of contents](#) | [back to top](#) ]

## **Program Information**


### **Long Term Ecological Research network (LTER)**

**Website:** <http://www.lternet.edu/>

**Coverage:** United States

**adapted from** <http://www.lternet.edu/>

The National Science Foundation established the LTER program in 1980 to support research on long-term ecological phenomena in the United States. The Long Term Ecological Research (LTER) Network is a collaborative effort involving more than 1800 scientists and students investigating ecological processes over long temporal and broad spatial scales. The LTER Network promotes synthesis and comparative research across sites and ecosystems and among other related national and international research programs. The LTER research sites represent diverse ecosystems with emphasis on different research themes, and cross-site communication, network publications, and research-planning activities are coordinated through the LTER Network Office.



- Site Codes**
- AND Andrews Forest LTER
  - ARC Arctic LTER
  - BES Baltimore Ecosystem Stu
  - BLE Beaufort Lagoon Ecosystems LTER
  - BNZ Bonanza Creek LTER
  - CCE California Current Ecosystem LTER
  - CDR Cedar Creek Ecosystem Science Reserve
  - CAP Central Arizona-Phoenix LTER
  - CWT Coweeta LTER
  - FCE Florida Coastal Everglades LTER
  - GCE Georgia Coastal Ecosystems LTER
  - HFR Harvard Forest LTER
  - HBR Hubbard Brook LTER
  - JRN Jornada Basin LTER
  - KBS Kellogg Biological Station LTER
  - KNZ Konza Prairie LTER
  - LUQ Luquillo LTER
  - MCM McMurdo Dry Valleys LT
  - MCR Moorea Coral Reef LTER
  - NWT Niwot Ridge LTER
  - NTL North Temperate Lakes I
  - NES Northeast U.S. Shelf LTER
  - NGA Northern Gulf of Alaska I
  - PAL Palmer Antarctica LTER
  - PIE Plum Island Ecosystems LTER
  - SBC Santa Barbara Coastal L
  - SEV Sevilleta LTER
  - VCR Virginia Coast Reserve L

2017 LTER research site map obtained from <https://lternet.edu/site/lter-network/>

[ [table of contents](#) | [back to top](#) ]

### Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1655686</a>
<a href="#">NSF Office of Integrative Activities (NSF OIA)</a>	<a href="#">OIA-1655221</a>
National Oceanic and Atmospheric Administration (NOAA)	<a href="#">NA18OAR4170094</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-2322676</a>

[ [table of contents](#) | [back to top](#) ]